

Nuclearity, Local Quasiequivalence and Split Property for Dirac Quantum Fields in Curved Spacetime

Claudio D'Antoni

*Dipartimento di Matematica, Università di Roma "Tor Vergata",
Via della Ricerca Scientifica,
I-00133 Roma, Italy*

Stefan Hollands

*Department of Physics, Enrico Fermi Institute,
University of Chicago, 5640 Ellis Ave.,
Chicago, IL 60367, U.S.A.*

February 6, 2008

Abstract

We show that a free Dirac quantum field on a globally hyperbolic spacetime has the following structural properties: (a) any two quasifree Hadamard states on the algebra of free Dirac fields are locally quasiequivalent; (b) the split-property holds in the representation of any quasifree Hadamard state; (c) if the underlying spacetime is static, then the nuclearity condition is satisfied, that is, the free energy associated with a finitely extended subsystem ("box") has a linear dependence on the volume of the box and goes like $\propto T^{s+1}$ for large temperatures T , where $s+1$ is the number of dimensions of the spacetime.

I Introduction

In the *algebraic framework* of quantum field theory [10], one takes the point of view that quantum field theoretical models should ultimately be described purely in terms of their associated algebras of local observables. The basic mathematical structure in this approach is an assignment $\mathcal{O} \rightarrow \mathfrak{A}(\mathcal{O})$ of finite regions in spacetime with C^* -algebras $\mathfrak{A}(\mathcal{O})$, containing the observables in the theory that are localized in $\mathcal{O}$. This assignment is expected to have some general, model-independent structural features; namely, if $\mathcal{O}_1 \subset \mathcal{O}_2$, then $\mathfrak{A}(\mathcal{O}_1) \subset \mathfrak{A}(\mathcal{O}_2)$ and if $\mathcal{O}_1$ and $\mathcal{O}_2$ are spacelike separated, then the elements in

$\mathfrak{A}(\mathcal{O}_1)$ should commute with those in $\mathfrak{A}(\mathcal{O}_2)$. The above properties express in an abstract way the general features of locality and local commutativity of a quantum field theory.

A quantum state in the algebraic framework is a positive normalized linear functional $\omega : \mathfrak{A} \rightarrow \mathbb{C}$. It is a basic fact from the theory of C^* -algebras that every state determines a representation π , called the GNS-representation, of $\mathfrak{A}$ on some Hilbert space $\mathcal{H}$ containing a distinguished vector $|\Omega\rangle$ such that $\omega(A) = \langle \Omega | \pi(A) \Omega \rangle$ for all $A \in \mathfrak{A}$. If the spacetime under consideration is Minkowski space, one furthermore postulates the existence of a preferred vacuum state, ω_0 , whose GNS-representation carries a positive energy representation of the translation symmetry group of Minkowski space.

The strength of the algebraic approach is that a number of structural features known from quantum field theory can be seen to be a consequence of general and model-independent axioms, see Haag's book [10] for an overview. However, these axioms do not yet entail such basic features as the existence of particles or a reasonable thermodynamic behavior. What is missing is a suitable mathematical formulation of the idea that excitations of the vacuum localized in finite regions and with finite energy occupy "finite volumes in phase space". Haag and Swieca [11] suggested to incorporate this idea into the general framework of algebraic quantum field theory in Minkowski space by demanding that a reasonable theory should satisfy the following *compactness criterion*: Let $\mathcal{O}_r$ be a double cone in Minkowski space whose base is a ball of radius r and let π_0 be the GNS-representation of the vacuum state on the Hilbert space $\mathcal{H}_0$, with vacuum vector $|\Omega_0\rangle$. Then the subsets of $\mathcal{H}_0$ given by

$$\{P_E \pi_0(A) |\Omega_0\rangle \in \mathcal{H}_0 \mid A \in \mathfrak{A}(\mathcal{O}_r), \|A\| \leq 1\}$$

are compact¹ for all E and r , where P_E is the spectral projector of the Hamiltonian, H , to the interval $[0, E]$.

With the same idea in mind, Buchholz and Wichmann [4] proposed a stronger requirement on the size of the phase space volumes corresponding to excitations localized in finite regions in Minkowski space and with finite energy. Instead of a sharp cut off in energy (represented above by the projector P_E) they use an exponential damping and replace the above compactness criterion by the following *nuclearity requirement*: The maps Θ_β defined by

$$\mathfrak{A}(\mathcal{O}_r) \ni A \rightarrow e^{-\beta H} \pi_0(A) |\Omega_0\rangle = \Theta_\beta(A) \in \mathcal{H}_0 \quad (1)$$

are nuclear for all $\beta > 0$, with *nuclearity index* bounded by $\nu_{r,\beta} \leq \exp[cr^s/\beta^n]$, where c, n are positive constants, s is the number of spatial dimensions and r the radius of the base of $\mathcal{O}_r$ (the definition of a nuclear map and its nuclearity index is given Sec. A.2). The physical interpretation of $\nu_{r,\beta}$ is that of the *partition function* of the finitely extended subsystem located in $\mathcal{O}_r$ at temperature $T = \beta^{-1}$. The dependence of $\nu_{r,\beta}$ on

¹Recall that a subset of a Banach space is called compact if every bounded sequence in this set contains a weakly convergent subsequence.

r then expresses that the free energy, $F_{r,\beta} = -\beta^{-1} \ln \nu_{r,\beta}$, of the finitely extended system depends at most linearly on the volume of the box. The dependence on β gives the relation between energy and temperature in the high temperature region. It has been shown to be sufficient to guarantee normal thermodynamic behavior.

The above nuclearity condition, as well as different variants thereof (see for example [5]), have been verified for free scalar fields and, more generally, for suitable infinite multiplets of these fields in Minkowski space [5]. A suitably modified nuclearity condition can also be formulated in curved spacetimes with a time translation symmetry, for which there exists a state ω_0 carrying a positive energy representation of the time translation group (so that H now denotes the generator of this symmetry). That the nuclearity property still holds under these more general circumstances was established by Verch [27] for the case of a free scalar field in the representation of the natural ground state on an ultrastatic spacetime.²

It is expected that the nuclearity property should hold also for free fields with higher spin, but a discussion of this seems to be missing from the literature so far, even in the case of Minkowski space. One purpose of the present paper is to fill this gap for the spin- $\frac{1}{2}$ case. Namely, we will prove that the nuclearity condition holds for the free Dirac field on a static spacetime in the GNS-representation of the ground state, and that the nuclearity index, $\nu_{\mathcal{C},\beta}$, associated with a double cone $\mathcal{O}_{\mathcal{C}}$ with compact base $\mathcal{C}$, satisfies the estimate

$$\nu_{\mathcal{C},\beta} \leq \exp \left[\left(\frac{\beta_0}{\beta} \right)^s e^{-m_0\beta/2} \right]. \quad (2)$$

Here, β_0 is a constant depending on the spatial geometry of the spacetime under consideration and m_0 is a constant proportional to the mass of the field. This estimate implies that the free energy $F_{\mathcal{C},\beta} = -\beta^{-1} \ln \nu_{\mathcal{C},\beta}$ associated with the subsystem localized in $\mathcal{O}_{\mathcal{C}}$ goes at most like T^{s+1} for high temperatures T . We also show that if the spacetime metric is rescaled by $g_{ab} \rightarrow \lambda^2 g_{ab}$, then the nuclearity index for the rescaled spacetime is estimated by the expression on the right side of (2), with β_0 replaced by $\lambda\beta_0$. This means that the free energy $F_{\mathcal{C},\beta}$ has at most a linear dependence on λ^s under rescalings of the metric, and therefore at most a linear dependence on the volume of the box.³

The proof of the above statements does not readily follow from the arguments developed for scalar fields, because the underlying combinatorics in both cases are quite different, due to the different statistics of bosonic and fermionic fields. Also, the above estimate for $\nu_{\mathcal{C},\beta}$ cannot readily be obtained by the same methods as in Minkowski space. Instead, we obtain it using some results and methods from the theory of pseudo differential operators. We mention as an aside that our proof of the nuclearity property also shows that the map (1) is actually even p -nuclear for all $p > 0$.

²Actually he verified a version of nuclearity condition, which is stronger than the one given above.

³Note that, in Minkowski space, the transformation $g_{ab} \rightarrow \lambda^2 g_{ab}$ is equivalent, via the conformal isometry $x \rightarrow \lambda x$, to a rescaling of the box itself, $\mathcal{O}_r \rightarrow \mathcal{O}_{\lambda r}$.

A quantum field model on a curved spacetime M is said to satisfy the so-called *split-property* (with respect to some state, ω) if the following holds: Let $\mathcal{O}_1$ and $\mathcal{O}_2$ be two concentric open double cones with bases $\mathcal{C}_1$ and $\mathcal{C}_2$ (contained in some Cauchy-surface Σ) such that $\text{clo}(\mathcal{C}_1) \subset \mathcal{C}_2$. Furthermore, let $\pi_\omega(\mathfrak{A}(\mathcal{O}_1))''$ and $\pi_\omega(\mathfrak{A}(\mathcal{O}_2))''$ be the weak closures of the corresponding local algebras in the GNS-representation of the state ω . Then there exists a type I factor $\mathcal{N}$ such that

$$\pi_\omega(\mathfrak{A}(\mathcal{O}_1))'' \subset \mathcal{N} \subset \pi_\omega(\mathfrak{A}(\mathcal{O}_2))''.$$

It is known [2] that the split-property automatically holds for the ground state ω_0 on an ultrastatic spacetime if the nuclearity condition is satisfied. Hence our estimate (2) immediately gives us that the split-property holds for free Dirac fields in static spacetimes in the GNS-representation of the ground state. (For the special case of a Dirac field in Minkowski spacetime this has previously been shown by Summers [26].) Combining this with a deformation argument due to Verch [27], we moreover show in Sec. V that the split-property holds in fact on an arbitrary globally hyperbolic spacetime in the GNS-representation of any quasifree Hadamard state (an explanation of this term will be given below). Our proof of this partly relies on the fact that any two quasifree Hadamard states ω_1 and ω_2 for the Dirac field are *locally quasiequivalent*, meaning that the density matrix states in the GNS Hilbert spaces of ω_1 and ω_2 define the same sets of states of $\mathfrak{A}(\mathcal{O})$ for any relatively compact subset $\mathcal{O}$ of M . The analogue of this result for scalar fields had previously been obtained by Verch [28], but a proof for Dirac case seems to be missing from the literature so far.⁴ It is therefore provided in Sec. IV. In the course of this proof we also show that the local v. Neumann algebras $\pi_\omega(\mathfrak{A}(\mathcal{O}_\mathcal{C}))''$ associated with a double cone $\mathcal{O}_\mathcal{C}$ (with regular base) is a *factor* for any quasifree state ω .

Because all quasifree Hadamard states are locally quasiequivalent, the v. Neumann algebras $\pi_\omega(\mathfrak{A}(\mathcal{O}))''$ are algebraically isomorphic for different choices of the quasifree state ω . They may therefore be regarded as different realizations of some abstract factorial v. Neumann algebra $\mathcal{R}(\mathcal{O})$. The collection of these factors can be shown to have the following *general covariance property* (for a detailed discussion of this property, see [29, 31]): Let $\chi : N \rightarrow M$ be a causality-preserving isometric embedding of a spacetime (N, h_{ab}) into a larger spacetime (M, g_{ab}) , so that $\chi^* g_{ab} = h_{ab}$, and assume that χ induces a homomorphism of the corresponding spin-structures. Then there exists a homomorphism α_χ such that $\alpha_\chi(\mathcal{R}(\mathcal{O})) = \mathcal{R}(\chi(\mathcal{O}))$ for all relatively compact $\mathcal{O} \subset N$.⁵ Moreover, if $\chi_1 \circ \chi_2$ is the composition of two isometric embeddings, then it holds that $\alpha_{\chi_1 \circ \chi_2} = \alpha_{\chi_1} \circ \alpha_{\chi_2}$. In the case when χ is a diffeomorphism, the map α_χ implements the transformation of local

⁴For scalar fields, it has recently been proven [16] that local quasiequivalence still holds under the much weaker assumption that the states in question are not Hadamard but only “adiabatic of order N ”, for some suitable number N . One would expect that an analogous result holds also for Dirac fields. A first result in this direction on Robertson-Walker spacetimes has been obtained by [13].

⁵As pointed out to us by the referee, in order that α_χ be independent of $\mathcal{O}$, one must actually require that the isometric embedding χ is extendible to some N' containing N as a subset with compact closure.

quantum fields under χ . Now if $\mathcal{O}_1 \subset \mathcal{O}_2 \subset N$ are open regions as in the discussion of the split-property, and if $\mathcal{N}$ is a type I factor corresponding to this inclusion, then $\alpha_\chi(\mathcal{N})$ is a type I factor corresponding to the inclusion $\chi(\mathcal{O}_1) \subset \chi(\mathcal{O}_2) \subset M$. In this sense, the type I factor $\mathcal{N}$ in the split-property ‘transforms in a generally covariant way’.

The plan and main results of this paper may now be summarized: In Section II, we review the quantization of a free Dirac field on a curved spacetime from the algebraic point of view and recall the notion of quasifree and Hadamard states. In Section III we show that any two quasifree Hadamard states are locally quasiequivalent (Thm. III.1), and that the local v. Neumann algebras associated with a quasifree state are factors of type III_1 (Thm. III.2). Section IV contains our results concerning the nuclearity property for Dirac fields in static spacetimes (Thm. IV.1 and Prop. IV.1). The split-property for Dirac fields in arbitrary globally hyperbolic spacetimes is established in the last section. The definition of nuclear maps and some related functional analytic concepts is given in the Appendix.

Notations and conventions: Throughout, (M, g_{ab}) denotes an oriented, time-oriented globally hyperbolic spacetime of dimension $s + 1$. The metric volume element compatible with the orientation of M is denoted by $\epsilon_{a_0 a_1 \dots a_s}$. We denote by $J^\pm(\mathcal{O})$ the causal future respectively past of a region $\mathcal{O} \subset M$, i.e., the set of all points that can be reached by a future respectively past directed causal curve starting in $\mathcal{O}$. $D(\mathcal{O})$ denotes the domain of dependence of $\mathcal{O}$, defined as the set of all points x , such any future and past inextendible causal curve through x intersects $\mathcal{O}$. Our signature convention is $+- - - \dots$.

II Review of classical and quantum Dirac fields in curved spacetimes

II.1 Classical fields

We begin by reviewing briefly the theory of a classical, linear Dirac field on a curved spacetime (for a more detailed discussion see for example [6, 23]). In order to define spinors on a curved spacetime, it is necessary to first introduce the notion of a spin-structure. Let FM be the bundle of all oriented, time-oriented orthonormal frames over (M, g_{ab}) . FM has the structure of a principal fibre bundle with base manifold M and structure group $SO^\uparrow(s, 1)$, where the arrow indicates that the transformations preserve the time-orientation. The universal 2–1 covering group of $SO^\uparrow(s, 1)$ is the group $\text{Spin}^\uparrow(s, 1)$. Elements g of this group can be represented by complex $2^{[(s+1)/2]}$ -dimensional⁶ square matrices satisfying

$$g\gamma^\mu g^{-1} = \Lambda(g)^\mu_\nu \gamma^\nu. \quad (3)$$

⁶ $[\dots]$ denotes the integer part of a number.

Here, the γ^μ denote a set of $2^{[(s+1)/2]}$ -dimensional gamma-matrices for $(s+1)$ -dimensional Minkowski spacetime⁷ satisfying

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2\eta^{\mu\nu}, \quad (4)$$

and Λ denotes the covering homomorphism. A spin structure on (M, g_{ab}) is by definition a principal fibre bundle SM with structure group $\text{Spin}^\uparrow(s, 1)$ and base manifold M , together with a bundle homomorphism $SM \rightarrow FM$ compatible with the covering homomorphism Λ . Whether a given spacetime admits a spin structure or not depends only on the topology of M . In the following we assume that M is such that a spin structure exists. Given a spin structure, one defines the “spinor bundle”, DM , as the $2^{[(s+1)/2]}$ -dimensional complex vector bundle over M which is associated to SM via the representation of $\text{Spin}^\uparrow(s, 1)$ on $\mathbb{C}^{2^{[(s+1)/2]}}$ defined by (3).

In order to write down the Dirac equation in M , one needs an analogue of Gamma-matrices as well as a suitable covariant derivative operator for spinor fields in curved spacetime. Gamma matrices γ^a in curved spacetime are the fiber-wise linear maps in DM satisfying $\gamma_a \gamma_b + \gamma_b \gamma_a = 2g_{ab}$. They are defined by requiring that their components with respect to an appropriate local frame are equal to those of the corresponding gamma matrices (3) in Minkowski space. Derivatives of spinor fields in curved spacetime can naturally be defined in terms of the so-called “spin-connection”, ∇_a , which is the uniquely determined covariant derivative operator satisfying $\nabla_a \gamma_b = 0$. The Dirac equation for a spinor field u in curved spacetime then reads

$$(i\gamma^a \nabla_a - m)u = 0, \quad m \geq 0. \quad (5)$$

It is known [6] that there exist unique advanced and retarded fundamental solutions $\mathcal{S}^{\text{adv}}$ and $\mathcal{S}^{\text{ret}}$ to this equation, i.e., continuous linear maps from $\mathcal{D}(DM)$ (the space of smooth spinor fields with compact support) to $\mathcal{E}(DM)$ (the space of smooth spinor fields), for which there holds

$$(i\gamma^a \nabla_a - m)\mathcal{S}^{\text{adv}} = \mathcal{S}^{\text{adv}}(i\gamma^a \nabla_a - m) = \mathbb{1},$$

(with the same equation holding for $\mathcal{S}^{\text{ret}}$) and for which $\text{supp}(\mathcal{S}^{\text{adv}}u) \subset J^+(\text{supp}(u))$ and $\text{supp}(\mathcal{S}^{\text{ret}}u) \subset J^-(\text{supp}(u))$. The causal propagator, $\mathcal{S}$, is the distributional bisolution defined by $\mathcal{S} = \mathcal{S}^{\text{adv}} - \mathcal{S}^{\text{ret}}$.

The notion of the charge-conjugate and the Dirac-conjugate of a spinor field u in Minkowski spacetime can be generalized in a natural and invariant way to spinor fields on a curved spacetime via an appropriately chosen trivialization of DM (the charge conjugation is defined only for $s+1 = 3, 4, 9, 10 \bmod 8$). The charge conjugate of a spinor field u is written as Cu , where C is a certain anti-linear, fiber-preserving, involutive ($C^2 = \mathbb{1}$) map

⁷We assume that the gamma matrices in (3) are taken in the Majorana representation, which requires that $s+1 = 3, 4, 9, 10 \bmod 8$.

from DM to itself. If u is a solution to the Dirac equation, then so is Cu . Spinor fields which are invariant under the charge conjugation map are called “Majorana spinors”. The Dirac-conjugate of a spinor field, $\bar{u}$, lives in the dual, D^*M , of the spinor bundle. If u is a solution to the Dirac equation (5), then $\bar{u}$ is a solution to $\bar{u}(-i\gamma^a\nabla_a - m) = 0$, and vice versa.

II.2 Quantum fields

The theory of a quantized Dirac field can be formulated in various ways. We here present the theory from the algebraic point of view, using Araki’s selfdual framework [1]. For simplicity, we will restrict attention from now on to Majorana fields (which means that $s+1 = 3, 4, 9, 10 \bmod 8$), but our results can easily be generalized to charged Dirac fields.

The algebra $\mathfrak{F}$ of Dirac-Majorana quantum fields is by definition the uniquely determined, unital C^* -algebra generated by elements of the form $\Psi(u)$ (smeared Dirac-Majorana quantum fields), where u is a compactly supported, smooth spinor field. They are subject to the following relations:

Linearity: The assignment $\mathcal{D}(DM) \ni u \rightarrow \psi(u) \in \mathfrak{F}$ is complex linear.

Dirac Equation: There holds $\Psi((i\gamma^a\nabla_a - m)u) = 0$.

Canonical Anticommutation Relations: $\{\Psi(u_1), \Psi(u_2)\} = i\mathcal{S}(C\bar{u}_1, u_2)\mathbb{1}$, where $\mathcal{S}$ is viewed as a bilinear form on $\mathcal{D}(D^*M) \times \mathcal{D}(DM)$.

Hermiticity: $\Psi(u)^* = \Psi(Cu)$.

The local field algebras $\mathfrak{F}(\mathcal{O})$ corresponding to some region $\mathcal{O}$ in spacetime, are by definition the C^* -subalgebras of $\mathfrak{F}$ generated by all elements of the form $\Psi(u)$, where $\text{supp}(u) \subset \mathcal{O}$.

In the following we will often make use of an alternative description of the algebra $\mathfrak{F}$ in terms of the Cauchy data of the field operators on some Cauchy surface Σ (“sharp time fields”), as we now briefly explain. It can be seen that

$$(k_1|k_2) = \int_{\Sigma} \bar{k}_1 \gamma^a k_2 \epsilon_{ab_1\dots b_s} \quad (6)$$

defines a positive, non-degenerate Hermitian inner product on the space of all compactly supported smooth spinor fields on Σ . The completion of this space with respect to this inner product is a Hilbert space which we will denote as $\mathcal{K} = L^2(DM \upharpoonright \Sigma)$. The charge conjugation map C can be seen to be antiunitary with respect to that inner product, that is, $(Ck_1|Ck_2) = (k_2|k_1)$ for all $k_1, k_2 \in \mathcal{K}$. One can show that $\mathfrak{F}$ is $*$ -isomorphic to the C^* -algebra generated by elements of the form $\psi(k)$ with $k \in \mathcal{K}$, subject to the relations $\psi(Ck) = \psi(k)^*$ and $\{\psi(k_1), \psi(k_2)\} = (Ck_1|k_2)\mathbb{1}$. The isomorphism is explicitly given by

$$\Psi(u) \rightarrow \psi(k), \quad k = \mathcal{S}u \upharpoonright \Sigma, \quad (7)$$

for spinor fields $u \in \mathcal{D}(DM)$. The local field algebras $\mathfrak{F}(\mathcal{O})$, where $\mathcal{O}$ is a region of the form $D(\mathcal{C})$ ($\mathcal{C}$ an open subset of Σ) correspond, under this isomorphism, to the algebras generated by elements of the form $\psi(k)$, where $k \in L^2(DM \upharpoonright \mathcal{C})$.

The algebra $\mathfrak{F}$ has the following structural property [1]: Any unitary operator U on $\mathcal{H}$ which commutes with C induces a unique automorphism α on $\mathfrak{F}$ satisfying $\alpha(\psi(k)) = \psi(Uk)$ for all $k \in \mathcal{H}$. α is called the “Bogoliubov automorphism” associated with U .

A state in the algebraic framework is by definition a linear functional $\omega : \mathfrak{F} \rightarrow \mathbb{C}$, which is normalized so that $\omega(\mathbb{1}) = 1$ and positive in the sense that $\omega(F^*F) \geq 0$ for all $F \in \mathfrak{F}$. The algebraic notion of states is related to the usual Hilbert space notion of states by the GNS-theorem. This says that for any algebraic state, there is a representation π_ω of $\mathfrak{F}$ as bounded, linear operators on a Hilbert space $\mathcal{H}_\omega$ containing a distinguished vector, $|\Omega_\omega\rangle$, such that $\omega(F) = \langle \Omega_\omega | \pi_\omega(F) \Omega_\omega \rangle$ for all $F \in \mathfrak{F}$. The two-point function, $\mathcal{S}_\omega$, of a state ω on $\mathfrak{F}$ is the bidistribution on $M \times M$ defined by

$$\mathcal{S}_\omega(u_1, u_2) = \omega(\Psi(u_1)\Psi(u_2))$$

for all smooth spinor fields u_1 and u_2 over M .

In this paper, we will restrict attention to a particular class of states on $\mathfrak{F}$, namely the so-called “quasifree states”. A quasifree state is one for which⁸

$$\omega(\Psi(u_1) \dots \Psi(u_n)) = \begin{cases} \sum_p \text{sign}(p) \prod_{(i,j) \in p} \omega(\Psi(u_i)\Psi(u_j)), & \text{for } n \text{ even,} \\ 0 & \text{for } n \text{ odd,} \end{cases} \quad (8)$$

where the sum is over all partitions p of $\{1, \dots, n\}$ into pairs (i, j) with $i < j$, and where $\text{sign}(p)$ is the signature of the permutation $\begin{pmatrix} n & n-1 & \dots & 2 & 1 \\ j_1 & i_1 & \dots & j_{n/2} & i_{n/2} \end{pmatrix}$. The above formula implies in particular that a quasifree state is entirely specified by its two-point function.

It is not difficult to see that the two-point function of a quasifree state can always be written in the form

$$\mathcal{S}_\omega(u_1, u_2) = \left(C\rho_\Sigma \mathcal{S}u_1 \middle| P\rho_\Sigma \mathcal{S}u_2 \right) \quad \text{for all } u_1, u_2 \in \mathcal{D}(DM), \quad (9)$$

where ρ_Σ means the operation of restricting a spinor field over M to the Cauchy surface Σ , $\mathcal{S}$ is the causal propagator and where P is a bounded operator on $\mathcal{H}$ satisfying $0 \leq P \leq \mathbb{1}$ and $CPC = \mathbb{1} - P$. In terms of the sharp time fields $\psi(k)$ (see eq. (7)), the two-point function is simply

$$\omega(\psi(k_1)\psi(k_2)) = (Ck_1 | Pk_2). \quad (10)$$

Conversely, given any operator P on $\mathcal{H}$ with the above properties, eq. (10) defines a quasifree state on $\mathfrak{F}$. If P is in addition a projection operator, $P^2 = P$, then the corresponding quasifree state is called a “Fock state”. A quasifree state is pure if and only if it

⁸An equivalent way of saying that a state is quasifree is to demand that it has a vanishing one-point function, $\omega(\psi(u)) = 0$, and vanishing truncated n -point functions for $n > 2$.

is a Fock state. The GNS representation of $\mathfrak{F}$ associated with a quasifree state is described as follows: $\mathcal{F}_\omega$ is the antisymmetric Fock-space over the “1-particle Hilbert space”

$$\mathcal{H} = \text{clo}\{P^{1/2}k \mid k \in C_0^\infty(DM \upharpoonright \Sigma)\}, \quad (11)$$

that is,

$$\mathcal{F}_\omega = \mathbb{C} \oplus \left(\bigoplus_{n \geq 1} (\wedge^n \mathcal{H}) \right), \quad (12)$$

where $\wedge^n \mathcal{H}$ is the n -th antisymmetrized tensor power of $\mathcal{H}$. The Hilbert space $\mathcal{F}_\omega$ is thus the closure of the linear span of vectors of the form

$$|\Phi\rangle = q_1 \wedge q_2 \wedge \cdots \wedge q_n, \quad q_i \in \mathcal{H}, \quad (13)$$

and the “vacuum vector”, $|\Omega_\omega\rangle$, corresponding to the first component $\mathbb{C}$ in the direct sum (12). The scalar product in $\mathcal{F}_\omega$ is that naturally induced from the scalar product $(\mid)$ in $\mathcal{H}$. The GNS-representation is

$$\pi_\omega(\psi(k)) = a(P^{1/2}k)^* + a(P^{1/2}Ck) \quad \forall k \in \mathcal{H}, \quad (14)$$

where $a(p)^*$ and $a(p)$ with $p \in \mathcal{H}$ are the creation respectively annihilation operators defined by⁹

$$\begin{aligned} a(p)^*(q_1 \wedge q_2 \wedge \cdots \wedge q_n) &= p \wedge q_1 \wedge q_2 \wedge \cdots \wedge q_n, \\ a(p)(q_1 \wedge q_2 \wedge \cdots \wedge q_n) &= \sum_{r=1}^n (-1)^{r+1} (p|q_r) q_1 \wedge \cdots \widehat{q_r} \wedge \cdots q_n, \quad q_i, p \in \mathcal{H}, \end{aligned} \quad (15)$$

as well as the action

$$a(p)^*|\Omega_\omega\rangle = p, \quad a(p)|\Omega_\omega\rangle = 0 \quad (16)$$

on the vacuum vector. They satisfy the usual anti-commutation relations

$$\{a(p_1), a(p_2)^*\} = (p_1|p_2). \quad (17)$$

Since the generators $\psi(k)$ of the algebra $\mathfrak{F}$ satisfy anticommutation relations, one has a slightly different notion of locality as compared to the case of Bose fields. In order to describe this notion of locality, it is convenient to introduce the grading automorphism γ on $\mathfrak{F}$, which is defined by $\gamma(\psi(k)) = -\psi(k)$ in terms of the generators of $\mathfrak{F}$. We have that $\gamma(F) = \pm F$ for $F \in \mathfrak{F}_\pm$, where $\mathfrak{F}_+$ is generated by even products of ψ , and where $\mathfrak{F}_-$ is generated by odd products of ψ . Each F can be uniquely decomposed into parts from $\mathfrak{F}_+$ and $\mathfrak{F}_-$ as $F = F_+ + F_-$, where $F_\pm = \frac{1}{2}(F \pm \gamma(F))$. Given this decomposition, one can introduce the graded commutator by

$$[F_1, F_2]_\gamma = \begin{cases} F_1 F_2 + F_2 F_1 & \text{if } F_1, F_2 \in \mathfrak{F}_-, \\ F_1 F_2 - F_2 F_1 & \text{otherwise,} \end{cases} \quad (18)$$

⁹Note that $p \rightarrow a(p)^*$ is linear in p and that $p \rightarrow a(p)$ is antilinear in p .

where we are assuming without loss of generality that $F_i \in \mathfrak{F}_\pm$. It is a consequence of the canonical anticommutation relations, together with locality properties of the anti-commutator function $\mathcal{S}$ that

$$[F_1, F_2]_\gamma = 0 \quad \text{if } F_1 \in \mathfrak{F}(\mathcal{O}_1), F_2 \in \mathfrak{F}(\mathcal{O}_2), \quad (19)$$

and if $\mathcal{O}_1$ and $\mathcal{O}_2$ are spacelike separated. The even part $\mathfrak{F}_+(\mathcal{O})$, but not the odd part, is a subalgebra of $\mathfrak{F}(\mathcal{O})$. Since the graded commutator is just the ordinary commutator for elements in the even part, it follows that

$$[F_1, F_2] = 0 \quad \text{if } F_1 \in \mathfrak{F}_+(\mathcal{O}_1), F_2 \in \mathfrak{F}_+(\mathcal{O}_2), \quad (20)$$

whenever $\mathcal{O}_1$ and $\mathcal{O}_2$ are spacelike separated. The algebra $\mathfrak{F}_+(\mathcal{O})$ therefore satisfies bosonic commutation relations appropriate for true observables in the theory. We will refer to it as the “algebras of observables”, and we will sometimes use the notation $\mathfrak{A}(\mathcal{O})$ for this algebra.

Actually, one can get rid of the graded commutator in eq. (18) in favor of the ordinary commutator via a standard construction often referred to as “twist”, and this leads to a slightly more subtle notion of locality, called “twisted locality”, for the algebra of fields, $\mathfrak{F}(\mathcal{O})$. We now briefly recall this concept, since it will play a role below. Any quasifree state ω is invariant under the automorphism γ , that is, $\omega(\gamma(F)) = \omega(F)$ for all $F \in \mathfrak{F}$, because such a state by definition vanishes on odd elements F . This implies that γ can be unitarily implemented in the GNS-representation of any quasifree state. In other words, there is a unitary U on $\mathcal{F}_\omega$ such that

$$U|\Omega_\omega\rangle = |\Omega_\omega\rangle, \quad \pi_\omega(\gamma(F)) = U\pi_\omega(F)U^* \quad \forall F \in \mathfrak{F}. \quad (21)$$

One can then define the “twisted (local) algebras”, $\pi_\omega(\mathfrak{F}(\mathcal{O}))^t$, by

$$\pi_\omega(\mathfrak{F}(\mathcal{O}))^t = \{\pi_\omega(F_+) + i\pi_\omega(F_-)U \mid F \in \mathfrak{F}(\mathcal{O})\}. \quad (22)$$

The twisted algebra $\pi_\omega(\mathfrak{F}(\mathcal{O}))^t$ is spatially isomorphic to $\pi_\omega(\mathfrak{F}(\mathcal{O}))$ and the isomorphism is implemented by the unitary $V = \frac{1}{\sqrt{2}}(1 + iU)$

In terms of the twisted algebras, one has the following modified version of locality, called “twisted locality”:

$$\pi_\omega(\mathfrak{F}(\mathcal{O}')) \subset \pi_\omega(\mathfrak{F}(\mathcal{O}))'^t. \quad (23)$$

Here, $\pi_\omega(\mathfrak{F}(\mathcal{O}))'$ denotes the commutant, that is, the set of all operators on $\mathcal{F}_\omega$ that commute with all $\pi_\omega(F)$, $F \in \mathfrak{F}(\mathcal{O})$. The set $\mathcal{O}'$ is the causal complement of the set $\mathcal{O}$, defined as the set of all spacetime points x such that $J(x) \cap \mathcal{O} = \emptyset$.

The above construction of Fock states is appropriate to obtain “ground states” for static spacetimes (M, g_{ab}) , which is the situation that we are considering in Sec. IV. Recall that a spacetime is called static if it has an everywhere time-like Killing vectorfield

t^a , $\mathcal{L}_t g_{ab} = 2\nabla_{(a} t_{b)} = 0$, which is orthogonal to a family of Cauchy surfaces Σ , or, equivalently, which satisfies $t_{[a} \nabla_b t_{c]} = 0$. We also assume here that the Killing field is timelike everywhere in the sense that $v^2 = t^a t_a \geq v_0^2 > 0$ for some v_0 , and that the orbits of t^a are complete. The construction of the ground state of the Dirac field for such a spacetime is precisely as follows: Consider a Cauchy surface Σ orthogonal to t^a . Since the Dirac equation has a well-posed initial value formulation, we may associate with any given $k \in C_0^\infty(DM \upharpoonright \Sigma)$ a uniquely determined solution u of the Dirac equation such that $u \upharpoonright \Sigma = k$. The flow $\{F_t\}_{t \in \mathbb{R}}$ of t^a induces a flow on the spinor bundle DM (denoted by the same symbol), which commutes with the charge conjugation¹⁰ and the Dirac operator [6]. Therefore, $F_t^* u$ is again a solution to the Dirac equation with initial data $k_t = F_t^* u \upharpoonright \Sigma$. It is not difficult to see that $k_t = \mathbf{u}(t)k$ for some 1-parameter group of unitary maps $\{\mathbf{u}(t)\}_{t \in \mathbb{R}}$ on $\mathcal{K}$ satisfying $[\mathbf{u}(t), C] = 0$ for all $t \in \mathbb{R}$. Hence, there is a 1-parameter family of Bogoliubov automorphisms $\{\alpha_t\}_{t \in \mathbb{R}}$ on $\mathfrak{F}$, given by $\alpha_t(\psi(k)) = \psi(\mathbf{u}(t)k)$ for all $k \in \mathcal{K}$. It is known [25, Prop. 4.1] that the 1-parameter group $\{\mathbf{u}(t)\}_{t \in \mathbb{R}}$ is strongly continuous on $\mathcal{K}$. Thus, by Stone's theorem, we can write

$$\mathbf{u}(t) = e^{it\mathbf{h}} \quad (24)$$

for some self-adjoint generator $\mathbf{h}$ on $\mathcal{K}$. It can be seen the action of $\mathbf{h}$ on a $k \in C_0^\infty(DM \upharpoonright \Sigma)$ is given by the differential operator

$$\mathbf{h}k = (-it^c \gamma_c q^{ab} \gamma_a \nabla_b + t^c \gamma_c m)k, \quad (25)$$

where $q_{ab} = g_{ab} - t_a t_b / v^2$ is the induced metric on Σ . Let now P be the projector on the positive energy subspace of $\mathbf{h}$,

$$P = \int_0^\infty E_{\mathbf{h}}(d\lambda), \quad (26)$$

with $E_{\mathbf{h}}$ the spectral measure of $\mathbf{h}$. Since $C\mathbf{h}C = -\mathbf{h}$, the projector P satisfies $CPC = \mathbb{1} - P$, and therefore defines a Fock state ω on $\mathfrak{F}$. That Fock state is, in fact, the ground state associated with the notion of time translations defined by the Killing vector field t^a .

The group $\{\alpha_t\}_{t \in \mathbb{R}}$ is unitarily implemented in the GNS-representation associated with the state ω in the sense that

$$\pi_\omega(\alpha_t(F)) = U(t)\pi_\omega(F)U(t)^* \quad \text{for all } F \in \mathfrak{F}, t \in \mathbb{R}, \quad (27)$$

¹⁰That can be seen e.g. as follows: Since the orbits of t^a are complete and since (M, g_{ab}) is assumed globally hyperbolic, every point $x \in M$ can be written uniquely as $x = F_t(y)$, $y \in \Sigma, t \in \mathbb{R}$. Since $F_t^* g_{ab} = g_{ab}$, we may assume that the set of spin-frames over any given point $x = F_t(y)$ is *defined* to be the set of spin-frames over the point y , and that $D_x M$ is consequently defined to be $D_y M$. This provides the desired (trivial) flow on the spin-bundle induced by F_t . The charge conjugation acting in the fiber $D_x M$ is defined to be the charge conjugation in $D_y M$ and therefore (trivially) commutes with the flow F_t .

where $U(t) = e^{itH}$ and where H is given by the second quantization of the self adjoint operator (the “1-particle Hamiltonian”) $\mathbf{h}_+ = P\mathbf{h}P$. H is the generator of the symmetry t^a on the GNS Hilbert space associated with ω and may therefore be identified with the Hamiltonian of the system. The vacuum vector $|\Omega_\omega\rangle$ is invariant under $U(t)$, i.e. $H|\Omega_\omega\rangle = 0$, thereby justifying the terminology “ground state” for ω . In Minkowski space, the above ground state construction gives the usual vacuum representation.

II.3 Hadamard states

In the previous subsection, we have recalled the definition of the algebra of observables for a quantized Dirac field on a curved spacetime, and we have introduced the notion of quasifree states on this algebra. We also recalled the construction of the ground state in a static spacetime, which is a preferred element in this class. The question then arises how to characterize a class of physically reasonable states on non-static, general globally hyperbolic spacetimes. It has been suggested for a long time that this class should include the quasifree states that have a two-point function of *Hadamard form*. As was recently shown by Hollands and Ruan [14], this is a necessary and sufficient condition in order to be able to extend the action of a quasifree state to more singular quantities such as Wick powers of the free field (for example stress-energy tensor)—which are not already contained in $\mathfrak{F}$ —as well as to other quantities arising in a perturbatively defined interacting quantum field theory.¹¹ We will see in Sec. V that the ground state on an ultrastatic spacetime (that is, a static spacetime for which the timelike Killing field is in addition normalized, $t^a t_a = -1$) is of Hadamard form.

For the convenience of the reader, we now recall the definition of Hadamard states for the Majorana-Dirac field. Roughly speaking, these are states whose two-point function has the short distance singularity structure of a Hadamard fundamental solution of the Dirac equation with no spacelike singularities. A mathematically precise definition of Hadamard states was first given by Kay and Wald [17] for the case of a scalar field and that definition has subsequently been generalized to Dirac fields by [30]. We here prefer to work with an alternative equivalent characterization of Hadamard states in terms of the so-called “wave front set” of their associated two-point function (for a definition of this concept see [12]). This characterization was first found by Radzikowski [21] in the case of scalar fields, and was later also established for Dirac fields, see [13, 18, 23].

¹¹More precisely in [14], the following general result was proven for a Hermitian scalar field: If the algebra $\mathfrak{F}$ of free quantum fields is enlarged to a suitable algebra $\mathfrak{W}$ containing the observables of interest in the perturbatively defined interacting theory, then the states on $\mathfrak{F}$ that can be extended to $\mathfrak{W}$ are precisely the Hadamard states with smooth truncated n -point functions for $n \neq 0$. This class includes in particular the quasifree Hadamard states, since these have vanishing truncated n -point functions for $n \neq 0$. The generalization of this result to Dirac fields is straightforward.

Definition II.1. A state ω on $\mathfrak{F}$ is said to be of Hadamard form if the wave front set, $\text{WF}(\mathcal{S}_\omega)$, of its two-point function is contained in the set

$$C^+ = \{(x_1, \xi_1; x_2, -\xi_2) \in T^*M^2 \setminus \{0\} \mid (x_1, \xi_1) \sim (x_2, \xi_2), \xi_1 \triangleright 0\}. \quad (28)$$

Here, the notation $(x_1, \xi_1) \sim (x_2, \xi_2)$ means that (a) the points x_1 and x_2 can be joined by a null-geodesic $c : [0, 1] \rightarrow M$, (b) the covectors ξ_1 and ξ_2 are cotangent to c (meaning that $\dot{c}^a(0) = g^{ab}(x_1)\xi_{1b}$ and $\dot{c}^a(1) = g^{ab}(x_2)\xi_{2b}$) and parallel transports of each other. $\xi_1 \triangleright 0$ means that ξ_1 is future-pointing.

Remarks. 1) An important consequence of the above definition is that if ω_1 and ω_2 are two Hadamard states, then the distribution $\mathcal{K} = \mathcal{S}_{\omega_1} - \mathcal{S}_{\omega_2}$ is given by a smooth function. To see this, we notice that

$$\mathcal{K}(Cu_1, Cu_2) = -\overline{\mathcal{K}(u_1, u_2)}, \quad (29)$$

by the Hermiticity and the commutation relations of the Dirac field. Since C is anti-linear, this equation implies, together with the definition of the wave front set, that if $(x_1, \xi_1; x_2, \xi_2) \in \text{WF}(\mathcal{K})$, then also $(x_1, -\xi_1; x_2, -\xi_2) \in \text{WF}(\mathcal{K})$; in other words

$$\text{WF}(\mathcal{K}) = -\text{WF}(\mathcal{K}). \quad (30)$$

On the other hand, we know that

$$\text{WF}(\mathcal{K}) = \text{WF}(\mathcal{S}_{\omega_1} - \mathcal{S}_{\omega_2}) \subset \text{WF}(\mathcal{S}_{\omega_1}) \cup \text{WF}(\mathcal{S}_{\omega_2}) \subset C^+, \quad (31)$$

because ω_1 and ω_2 are Hadamard. Since the only subset of $S \subset C^+$ with $S = -S$ is the empty set, this implies that in fact $\text{WF}(\mathcal{K}) = \emptyset$, which is equivalent to the statement that $\mathcal{K}$ is smooth.

2) If a quasifree state is of Hadamard form in a globally hyperbolic neighbourhood of a Cauchy surface, then, as originally shown by Fulling, Narcowich and Wald [8], it must in fact be of Hadamard form throughout the whole spacetime. For a more recent version of this argument within the framework of microlocal analysis, see for example [13, 18, 23].

III Local quasiequivalence and factoriality

We will now show that the GNS representations π_{ω_1} and π_{ω_2} of any two quasifree Hadamard states ω_1 and ω_2 on $\mathfrak{F}$ are quasiequivalent when restricted to $\mathfrak{F}(\mathcal{O})$ for any open, relatively compact set $\mathcal{O} \subset M$, meaning that the density matrices in the GNS Hilbert spaces $\mathcal{F}_{\omega_1}$ and $\mathcal{F}_{\omega_2}$ define the same sets of states of $\mathfrak{F}(\mathcal{O})$.

In order to prove this result, we will proceed as follows: We first note that $\pi_{\omega_1} \upharpoonright \mathfrak{F}(\mathcal{O})$ is quasiequivalent to $\pi_{\omega_2} \upharpoonright \mathfrak{F}(\mathcal{O})$, if $\pi_{\omega_1} \upharpoonright \mathfrak{F}(\tilde{\mathcal{O}})$ is quasiequivalent to $\pi_{\omega_2} \upharpoonright \mathfrak{F}(\tilde{\mathcal{O}})$ for some

$\tilde{\mathcal{O}} \supset \mathcal{O}$. Now every open, relatively compact set $\mathcal{O} \subset M$ can be embedded into a set of the form $\text{int}(D(\mathcal{C}))$, where $\mathcal{C}$ is a relatively compact subset of some Cauchy surface Σ with smooth boundary in Σ ¹². We may therefore restrict attention to regions $\mathcal{O}_{\mathcal{C}}$ of the form $\text{int}(D(\mathcal{C}))$. For such regions, we first show that the GNS representation $\pi_{\omega|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}}$ of the partial state $\omega|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ is quasiequivalent to the representation $\pi_{\omega}|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$. Then we will check that the representations relative to any partial (quasifree, Hadamard) state $\omega|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ are locally quasiequivalent.

Lemma III.1. Let $A \supset B$ be C^* -algebras, ω a state on A , and $(\pi, \mathcal{F}, |\Omega\rangle)$ be the GNS representation of A . Denote by e' the orthogonal projection on $\pi(B)|\Omega\rangle$. Then $\pi_{\omega|_B}$ is quasiequivalent to $\pi_{\omega}|_B$ if and only if the central support of e' is $\mathbb{1}$.

Proof. The projection e' belongs to $\pi(B)'$ and $B \ni b \rightarrow \pi(b)|_{e'\mathcal{F}}$ is equivalent to the GNS representation of $\omega|_B$. If the central support of e' in $\pi(B)'' \cap \pi(B)'$ is equal to $\mathbb{1}$, then the homomorphism (induction) $\pi(B)'' \ni T \rightarrow T|_{e'\mathcal{F}}$ is an isomorphism that gives the quasiequivalence between $B \ni b \rightarrow \pi(b)$ and $B \ni b \rightarrow \pi(b)|_{e'\mathcal{F}}$. Conversely, quasiequivalent representations have equivalent amplifications. $\square$

In order to be able to apply the above lemma in the case at hand we next prove a proposition that is also of some interest in its own right.

Proposition III.1. Let $\mathcal{C}$ be a relatively compact subset of a Cauchy surface Σ with smooth boundary in Σ , and let $\mathcal{O}_{\mathcal{C}} = \text{int}(D(\mathcal{C}))$, where $D(\mathcal{C})$ is the domain of dependence of $\mathcal{C}$. Then, for any quasifree state ω ,

$$\pi_{\omega}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))^t \cap \pi_{\omega}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))'' = \mathbb{C} \cdot \mathbb{1}. \quad (32)$$

Remark: Taking into account the even and odd parts $\mathfrak{F}_{\pm}$ separately, the proposition implies that the even part of the center of $\pi_{\omega}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))$ is trivial.

Proof. Let P be the operator on $\mathcal{H} = L^2(DM|_{\Sigma})$ corresponding to the quasifree state ω via eq. (9), and let $\mathcal{H} \subset \mathcal{H}$ be the 1-particle Hilbert space, as defined in eq. (11). In addition, let us define the following closed, real linear subspaces of $\mathcal{H}$:

$$\begin{aligned} \mathcal{M}(\mathcal{C}) &= \text{clo}\{P^{1/2}k \mid k \in C_0^{\infty}(DM|_{\mathcal{C}}), \quad Ck = k\}, \\ \mathcal{M} &= \text{clo}\{P^{1/2}k \mid k \in C_0^{\infty}(DM|_{\Sigma}), \quad Ck = k\}. \end{aligned}$$

¹²Proof: Let K be a compact set containing $\mathcal{O}$, and let $\mathcal{C} = (J^+(K) \cap \Sigma) \cup (J^-(K) \cap \Sigma)$. Since Σ is a Cauchy surface, it follows that $D(\mathcal{C}) \supset K$ by the definition of the domain of dependence. We will show that $\mathcal{C}$ is compact by showing that $J^{\pm}(K) \cap \Sigma$ is compact. The collection of sets $\{J^-(x)\}_{x \in M}$ is a cover of K and hence has a finite subcover $\{J^-(x_1), \dots, J^-(x_r)\}$. By Thm. 8.3.12 of [32], $J^-(x) \cap \Sigma$ is compact for any point x , and therefore so is also $\cup_{i=1}^r (J^-(x_i) \cap \Sigma)$. Hence, $J^-(K) \cap \Sigma$ is contained in a compact set. But $J^-(K) \cap \Sigma$ is also closed by Thm. 8.3.11 of [32], hence compact. The same argument can be repeated for “+”, showing that $\mathcal{C}$ is a compact subset of Σ . By enlarging $\mathcal{C}$ if necessary, we can achieve that it has a smooth boundary in Σ .

We note that $\mathcal{H} = \mathcal{M} + i\mathcal{M}$ and it is not difficult to see that $\mathcal{H} = \mathcal{M}$ if and only if P is a projector. By eq. (14), we can write

$$\pi_\omega(\mathfrak{F}(\mathcal{O}_\mathcal{C}))'' = \{a(m) + a(m)^* \mid m \in \mathcal{M}(\mathcal{C})\}''.$$

By an argument due to Foit [7] and Roberts, we know that (32) holds if and only if $\mathcal{M}(\mathcal{C}) \cap i\mathcal{M}(\mathcal{C})' = \{0\}$, where “prime” denotes the symplectic complement of a set in $\mathcal{H}$, defined as

$$\mathcal{M}(\mathcal{C})' = \{m' \in \mathcal{H} \mid \text{Im}(m|m') = 0 \quad \forall m \in \mathcal{M}(\mathcal{C})\}.$$
 (33)

Let now $m \in \mathcal{M}(\mathcal{C}) \cap i\mathcal{M}(\mathcal{C})'$. We must show that $m = 0$. Now, since $m \in i\mathcal{M}(\mathcal{C})'$, we have that $im \in \mathcal{M}(\mathcal{C})'$, which means that $\text{Re}(m|m') = 0$ for all $m' \in \mathcal{M}(\mathcal{C})$. Moreover, since $m \in \mathcal{M}(\mathcal{C})$, it is not difficult to see that $\text{Re}(m|m') = 0$ for all $m' \in \mathcal{M}(\Sigma \setminus \mathcal{C})$. Altogether, this means that

$$\text{Re}(m|m') = 0 \quad \forall m' \in \mathcal{M}(\mathcal{C}) + \mathcal{M}(\Sigma \setminus \mathcal{C}).$$
 (34)

Consequently, we know that

$$\text{Re}(m|P^{1/2}k') = 0 \quad \forall k' \in C_0^\infty(DM \upharpoonright \mathcal{C}) + C_0^\infty(DM \upharpoonright \Sigma \setminus \mathcal{C}), \quad Ck' = k'. \quad (35)$$

Because the boundary of $\mathcal{C}$ is smooth, it follows easily that the space $C_0^\infty(DM \upharpoonright \mathcal{C}) + C_0^\infty(DM \upharpoonright \Sigma \setminus \mathcal{C})$ is dense in $C_0^\infty(DM \upharpoonright \Sigma)$ in the norm induced by the scalar product (6). This implies that

$$\text{Re}(m|m') = 0 \quad \forall m' \in \mathcal{M}, \quad (36)$$

or equivalently that $m \in \mathcal{M} \cap i\mathcal{M}'$. Let us now introduce an auxiliary quasifree state $\tilde{\omega}$ whose corresponding operator $\tilde{P}$ is a projection, so that $\tilde{\mathcal{H}} = \tilde{\mathcal{M}}$ holds for the corresponding 1-particle space. Let us write $m = \lim_{j \rightarrow \infty} P^{1/2}k_j$, where $k_j \in C_0^\infty(DM \upharpoonright \Sigma)$ with $Ck_j = k_j$. Then, since

$$\|P^{1/2}k_i - P^{1/2}k_j\| = \|k_i - k_j\| = \|\tilde{P}^{1/2}k_i - \tilde{P}^{1/2}k_j\|, \quad (37)$$

it follows that the limit $\tilde{m} = \lim_{j \rightarrow \infty} \tilde{P}^{1/2}k_j$ exists in $\tilde{\mathcal{M}}$. If $m' = P^{1/2}k'$, with k' an arbitrary element of $C_0^\infty(DM \upharpoonright \Sigma)$ with $Ck' = k'$, then (36) implies that

$$\begin{aligned} 0 &= \text{Re}(m|m') \\ &= \lim_{j \rightarrow \infty} \text{Re}(P^{1/2}k_j|P^{1/2}k') \\ &= \lim_{j \rightarrow \infty} \text{Re}(\tilde{P}^{1/2}k_j|\tilde{P}^{1/2}k') \\ &= \text{Re}(\tilde{m}|\tilde{m}'), \end{aligned}$$

where $\tilde{m}' = \tilde{P}^{1/2}k'$. It follows from this that $\text{Re}(\tilde{m}|\tilde{m}')$ for all $\tilde{m}' \in \tilde{\mathcal{M}} = \tilde{\mathcal{H}}$, and therefore that $\tilde{m} = 0$. But, by (37), this also implies that $m = 0$, which proves the proposition. $\square$

We now will now combine the above two results to show

Proposition III.2. Let ω be a quasifree state on $\mathfrak{F}$. Then the representations $\pi_{\omega}|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ and $\pi_{\omega}|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ are quasiequivalent.

Proof. Let U be the unitary operator on $\mathcal{H}_{\omega}$ implementing the automorphism γ , $\pi_{\omega}(\gamma(F)) = U\pi_{\omega}(F)U^*$. By the remark following Prop. III.1, we know that the $\text{Ad}(U)$ -invariant (that is, even) part of $\pi_{\omega}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))'' \cap \pi_{\omega}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))'$ is trivial. The central support of $\pi_{\omega}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))|_{\Omega_{\omega}}$ is invariant under $\text{Ad}(U)$, hence it is $\mathbb{1}$. The result now follows from Lem. III.1. $\square$

Theorem III.1. (Local quasiequivalence) The GNS representations π_{ω_1} and π_{ω_2} of any two quasifree Hadamard states ω_1 and ω_2 are locally quasiequivalent when restricted to $\mathfrak{F}(\mathcal{O})$ for any open, relatively compact set $\mathcal{O} \subset M$.

Proof. As explained above, we only need to prove the quasiequivalence of the representations $\pi_{\omega_1}|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ and $\pi_{\omega_2}|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$, where $\mathcal{O}_{\mathcal{C}}$ is a region of the form $\text{int}(D(\mathcal{C}))$. Let P_1 and P_2 be the self-adjoint operators on $\mathcal{H} = L^2(DM|_{\Sigma})$ associated to the quasifree states ω_1 and ω_2 via eq. (9). The partial states $\omega_1|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ and $\omega_2|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ are then again quasifree states, corresponding to the operators $E_{\mathcal{C}}P_1E_{\mathcal{C}}$ and $E_{\mathcal{C}}P_2E_{\mathcal{C}}$, where $E_{\mathcal{C}}$ is the projector onto the closed subspace $L^2(DM|_{\mathcal{C}})$ of $\mathcal{H}$, given by $(E_{\mathcal{C}}k)(x) = \chi_{\mathcal{C}}(x)k(x)$ with $\chi_{\mathcal{C}}$ the characteristic function of the set $\mathcal{C}$.

By a theorem of Powers and Størmer [20, Thm. 5.1], the partial states $\omega_1|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ and $\omega_2|_{\mathfrak{F}(\mathcal{O}_{\mathcal{C}})}$ are guaranteed to be quasiequivalent if

$$\|(E_{\mathcal{C}}P_1E_{\mathcal{C}})^{1/2} - (E_{\mathcal{C}}P_2E_{\mathcal{C}})^{1/2}\|_2 < \infty, \quad (38)$$

where $\|\cdot\|_2$ is the Hilbert-Schmidt norm on $\mathcal{H}$, see Sec. A.2. In order to estimate the left side of (38), we recall the Powers-Størmer inequality, which says that $\|A^{1/2} - B^{1/2}\|_2^2 \leq \|A - B\|_1$ for any two positive bounded operators A and B on a Hilbert space, where $\|\cdot\|_1$ denotes the trace-norm, see Sec. A.2. Taking $E_{\mathcal{C}}P_1E_{\mathcal{C}}$ for A and $E_{\mathcal{C}}P_2E_{\mathcal{C}}$ for B in that inequality, we find that (38) follows if we can show that

$$\|E_{\mathcal{C}}(P_1 - P_2)E_{\mathcal{C}}\|_1 < \infty. \quad (39)$$

We now define a bidistribution K on $\mathcal{D}(DM|_{\Sigma}) \times \mathcal{D}(DM|_{\Sigma})$ associated with the operator $E_{\mathcal{C}}(P_1 - P_2)E_{\mathcal{C}}$ by putting $K(k_1, k_2) = (Ck_1|E_{\mathcal{C}}(P_1 - P_2)E_{\mathcal{C}}k_2)$. This bidistribution can be expressed in terms of the two-point functions $\mathcal{S}_{\omega_1}$ and $\mathcal{S}_{\omega_2}$ as follows: By the Hadamard property combined with a microlocal argument (see e.g. [13]), the latter possess a well-defined pull-back to the submanifold $\Sigma \times \Sigma$, denoted by $\varphi_{\Sigma}^*\mathcal{S}_{\omega_1}$ respectively $\varphi_{\Sigma}^*\mathcal{S}_{\omega_2}$, where $\varphi_{\Sigma} : \Sigma \times \Sigma \rightarrow M \times M$ is the embedding map. Using eq. (9), we compute that

$$\begin{aligned} \varphi_{\Sigma}^*\mathcal{S}_{\omega_j}(\gamma_a n^a k_1, \gamma_b n^b k_2) &= (C\rho_{\Sigma}\mathcal{S}'_{\Sigma}(\gamma_a n^a k_1)|P_j\rho_{\Sigma}\mathcal{S}'_{\Sigma}(\gamma_b n^b k_2)) \\ &= (Ck_1|P_jk_2) \quad (j = 1, 2), \end{aligned} \quad (40)$$

where n^a is the normal of Σ and where we have used the identity $\rho_\Sigma \mathcal{S} \rho'_\Sigma = i \gamma_a n^a \mathbb{1}_\Sigma$ (see [6]) in the last line, with ρ'_Σ the dual of the restriction map ρ_Σ . It is not difficult to see from this that

$$K(k_1, k_2) = \varphi_\Sigma^*(\mathcal{S}_{\omega_1} - \mathcal{S}_{\omega_2})(\chi_{\mathcal{C}} \gamma_a n^a k_1, \chi_{\mathcal{C}} \gamma_b n^b k_2) \quad (41)$$

Since ω_1 and ω_2 are by assumption Hadamard states, we know that $\mathcal{S}_{\omega_1} - \mathcal{S}_{\omega_2}$ is smooth on $M \times M$ (see the first remark following Def. II.1). Equation (41) therefore implies that K can be identified with a smooth section in $D^*M \upharpoonright \Sigma \times D^*M \upharpoonright \Sigma$, with compact support in $\mathcal{C} \times \mathcal{C}$. Let $\{\eta_i\}$ be a smooth partition of unity covering $\mathcal{C}$ with the property that the support of each η_i is contained in a coordinate patch. Then we can write $K = \sum (\eta_i \otimes \eta_j) K = \sum K_{ij}$, and each K_{ij} can be identified a matrix of smooth, compactly supported functions in $\mathcal{D}(\mathbb{R}^s \times \mathbb{R}^s)$ via a local trivialization of $D^*M \upharpoonright \Sigma \times D^*M \upharpoonright \Sigma$ over the patch corresponding to the pair η_i, η_j . As it is well known, a matrix of operators is in the trace class if each matrix entry is in the trace class. Thus, in order to show that K is in the trace class, it suffices to show that the matrix elements of each K_{ij} is in the trace class. In fact any smooth, compactly supported integral kernel L in $\mathcal{D}(\mathbb{R}^s \times \mathbb{R}^s)$ is in the trace class, as one can see e.g. as follows: We can write

$$L(x, y) = (1 + |x|)^{-p} (1 + |x|)^p L(x, y), \quad (42)$$

where $|x| = (\sum x_i^2)^{1/2}$. The multiplication operator by $(1 + |x|)^{-p}$ is in the Hilbert-Schmidt class for p sufficiently large since $\|(1 + |x|)^{-p}\|_2^2 = \int (1 + |x|)^{-2p} d^s x < \infty$ then, and the integral kernel $(1 + |x|)^p L(x, y)$ is in $L^2(\mathbb{R}^s \times \mathbb{R}^s)$ for any p , and therefore also in the Hilbert-Schmidt class. Hence, L is the product of two Hilbert-Schmidt operators, and hence in the trace class. $\square$

We conclude this section with the following theorem.

Theorem III.2. (Local factoriality) Let ω be a quasifree state Hadamard state for the free Dirac field on a globally hyperbolic spacetime. Then the local v. Neumann algebras $\pi_\omega(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))''$ are type III_1 factors. Furthermore, the local quasiequivalence established in Thm. III.1 is in fact local unitary equivalence.

Proof. The partial state $\omega \upharpoonright \mathfrak{F}(\mathcal{O}_{\mathcal{C}})$ is a quasifree state corresponding to $E_{\mathcal{C}} P E_{\mathcal{C}}$ in the notation of the proof of Thm. III.1. Hence, the algebra $\pi_{\omega \upharpoonright \mathfrak{F}(\mathcal{O}_{\mathcal{C}})}(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))$ is a factor by the results of Powers and Størmer [20]. By local quasiequivalence, also $\pi_\omega(\mathfrak{F}(\mathcal{O}_{\mathcal{C}}))$ is a factor. The type of this factor can be obtained by again invoking local quasiequivalence. In fact, the local v. Neumann algebras are type III_1 factors in a pure, quasifree Hadamard state by [30, Cor. 5.12], therefore they are III_1 factors for all quasifree Hadamard states. $\square$

Remark. By a similar argument, one can also show that the local algebras of observables, $\pi_\omega(\mathfrak{A}(\mathcal{O}_{\mathcal{C}}))''$ are factors of type III_1 .

IV Nuclearity

In this section, we show that the theory of a free Dirac field on a static spacetime satisfies the nuclearity condition stated in the introduction. Before giving the precise formulation of our result, let us begin by fixing our conventions for this section, as well as some notations: We assume that the spacetime (M, g_{ab}) under consideration is static. Let Σ_t be the Cauchy surfaces orthogonal to the timelike Killing vector field. We pick an arbitrary but fixed $\Sigma = \Sigma_t$, and fix a subset $\mathcal{C} \subset \Sigma$ with compact closure. For that subset, we define $\mathcal{O}_{\mathcal{C}} = \text{int}(D(\mathcal{C})) \subset M$, where $D(\mathcal{C})$ is the domain of dependence of $\mathcal{C}$. Throughout, we will work in the representation of $\mathfrak{F}$ given by the ground state ω introduced in Sec. II. In this representation, the Fermi fields take the form

$$\psi(k) = a(Pk)^* + a(PCk), \quad k \in \mathcal{K} = L^2(\Sigma; DM), \quad (43)$$

where we are writing $\psi(k)$ instead of $\pi_{\omega}(\psi(k))$ to lighten the notation in this section. Here, P is the projector on the positive spectral subspace of $\mathbf{h}$ (see eq. (26)), and $a(p)^*$ and $a(p)$ are the creation resp. annihilation operators defined in eq. (15), with $p \in \mathcal{H} = P\mathcal{K}$. The local algebra $\mathfrak{F}(\mathcal{O}_{\mathcal{C}})$ is then generated by those $\psi(k)$ for which k is supported in $\mathcal{C}$, that is $k \in L^2(\mathcal{C}; DM)$. It is viewed as a subalgebra of the algebra of bounded operators on $\mathcal{F}_{\omega}$ via π_{ω} . For further notational simplicity, we will also drop the subscript “ ω ” on $\mathcal{F}_{\omega}$ and the vacuum vector $|\Omega_{\omega}\rangle$ in this section, the understanding being that we always refer to the quantities associated with the ground state. For the relevant definitions concerning the nuclearity of a map and related functional analytic concepts we refer the reader to Sec. A.2.

Let Θ_{β} be the map defined by

$$\Theta_{\beta} : \mathfrak{F}(\mathcal{O}_{\mathcal{C}}) \rightarrow \mathcal{F}, \quad \Theta_{\beta}(A) = e^{-\beta H} A |\Omega\rangle, \quad (44)$$

for $\beta > 0$, and let

$$\nu_{\mathcal{C}, \beta} = \|\Theta_{\beta}\|_1$$

be its nuclearity index. The aim of the present section is to show that Θ_{β} is p -nuclear (Thm. IV.1) and to estimate the nuclearity index $\nu_{\mathcal{C}, \beta}$ (Prop. IV.1).

Theorem IV.1. (Nuclearity) The maps Θ_{β} are p -nuclear for all $\beta > 0$ and all $p > 0$, i.e., $\|\Theta_{\beta}\|_p < \infty$.

Proof: We first show that p -nuclearity of the maps Θ_{β} follows if, for each $\beta > 0$, there is a positive operator T on $\mathcal{H} = P\mathcal{K}$, with the following two properties:

- (a) $T \in \mathcal{I}_p(\mathcal{H})$ for all $p > 0$ and all $\beta > 0$, where $\mathcal{I}_p(\mathcal{H})$ denotes the Schatten-space with index p , consisting of all bounded operators B such that $\|B\|_p = (\text{tr}|B|^p)^{1/p} < \infty$ (see also Sec. A.2).

- (b) If $\{p_j\}_{j \in \mathbb{N}}$ is an orthonormal basis of $\mathcal{H}$ consisting of eigenvectors of T and $\{t_j\}_{j \in \mathbb{N}}$ the corresponding eigenvalues,

$$Tp_j = t_j p_j,$$

then there holds the inequality

$$|\langle \Phi_{i_1 \dots i_k} | \Theta_\beta(A) \rangle| \leq t_{i_1} \dots t_{i_k} \|A\| \quad \text{for all } A \in \mathfrak{F}(\mathcal{O}_\mathcal{C}), \quad (45)$$

where $|\Phi_{i_1 \dots i_k}\rangle = a(p_{i_1})^* \dots a(p_{i_k})^* |\Omega\rangle$ with $i_1 < \dots < i_k$.

To see that the existence of such an operator T indeed implies the statement of the theorem, we expand Θ_β as

$$\Theta_\beta(A) = \sum_{k \geq 0} \sum_{i_1 < \dots < i_k} |\Phi_{i_1 \dots i_k}\rangle \varphi_{i_1 \dots i_k}(A)$$

where $\varphi_{i_1 \dots i_k}(A) = \langle \Phi_{i_1 \dots i_k} | \Theta_\beta(A) \rangle$. Since $\| |\Phi_{i_1 \dots i_k}\rangle \| = 1$ and since $\|\varphi_{i_1 \dots i_k}\| \leq t_{i_1} \dots t_{i_k}$ by eq. (45) in assumption (b), we get the estimate

$$\begin{aligned} \|\Theta_\beta\|_p &\leq \left(\sum_{k \geq 0} \sum_{i_1 < \dots < i_k} \|\varphi_{i_1 \dots i_k}\|^p \right)^{\frac{1}{p}} \\ &\leq \left(\sum_{k \geq 0} \sum_{i_1 < \dots < i_k} t_{i_1}^p \dots t_{i_k}^p \right)^{\frac{1}{p}} \\ &= \left(\sum_{k \geq 0} \text{tr}(\wedge^k(T^p)) \right)^{\frac{1}{p}} = (\det(\mathbb{1} + T^p))^{\frac{1}{p}}, \end{aligned}$$

where in the last line we have used a well-known identity for trace-class operators, see e.g. [22, Thm. XIII.106]. Using the inequality [22, Lem. 4, Sec. XIII.17],

$$|\det(\mathbb{1} + T^p)| \leq e^{\|T^p\|_1},$$

we conclude that

$$\|\Theta_\beta\|_p \leq e^{\frac{1}{p} \|T\|_p^p} \quad (46)$$

which is finite, because $T \in \mathcal{I}_p(\mathcal{H})$ by assumption (a). This then shows that the maps Θ_β are p -nuclear for any $p > 0$ and $\beta > 0$, with $\|\Theta_\beta\|_p \leq (\det(\mathbb{1} + T^p))^{1/p}$.

In order to complete the proof, we must show the existence of an operator T with the above properties (a) and (b). Let $E_\mathcal{C}$ be the projector onto the closed subspace $L^2(DM \upharpoonright \mathcal{C})$ of $\mathcal{H}$ (i.e., multiplication by the characteristic function of the set $\mathcal{C}$), define

$$S = 2E_\mathcal{C} e^{-\beta \mathbf{h}} P, \quad (47)$$

and let $S = VT$ be the polar decomposition of S (so that $T = (S^*S)^{1/2}$). In what follows, we will establish that the so defined operator T has the above stated properties (a) and (b). Let us begin by stating some relevant properties of S and T .

Lemma IV.1. The operators T and S have the following properties:

- (i) $T = PT = TP$ and $\|T\| \leq \sqrt{2}e^{-\beta m_0}$.
- (ii) $T \in \mathcal{I}_p(\mathcal{H})$ for all $p > 0$ for all $\beta > 0$.
- (iii) There holds the estimate

$$\|S\|_1 \leq \left(\frac{\beta_0}{\beta}\right)^s e^{-\beta m_0/2},$$

where s is the number of spatial dimensions, $m_0 = v_0 m$ and $\beta_0 > 0$ is a constant independent of β .

- (iv) The range of S^* is dense in $\mathcal{H}$.

Proof. The first part of (i) follows immediately from the definition. In order to show the norm estimate in (i), we notice that (25) implies that

$$\mathbf{h}^2 \geq m_0^2 \mathbb{1}, \quad (48)$$

where $m_0 = v_0 m$. We therefore get the inequality $e^{-\beta \mathbf{h}} P \leq e^{-m_0 \beta}$ and hence $\|T\| = \|T^2\|^{1/2} = \sqrt{2} \|e^{-\beta \mathbf{h}} P E_{\mathcal{C}} P e^{-\beta \mathbf{h}}\|^{1/2} \leq \sqrt{2} e^{-\beta m_0}$. Part (ii) of the lemma can be demonstrated by adapting a method developed in [3], we briefly sketch the argument. Let $\chi \in C_0^\infty(\Sigma)$ be a function which is equal to one on $\mathcal{C}$ and let M_χ be the corresponding multiplication operator on $\mathcal{K} = L^2(DM \upharpoonright \Sigma)$, defined by $M_\chi k = \chi k$. Then one can show that the operators

$$\mathbf{k}_n = (\mathbb{1} + \beta^2 \mathbf{h}^2)^{(n-1)s/2} M_\chi (\mathbb{1} + \beta^2 \mathbf{h}^2)^{-ns/2}$$

are in the Hilbert-Schmidt class for all $n \geq 1$. A proof of this is given in the Appendix of ref. [4] for the case of Minkowski space; the main steps for a proof of this statement in the case when Σ is not flat are given in the Appendix of this paper. Now it trivially follows from the definition (47) of S that one can write

$$S = 2E_{\mathcal{C}} \cdot \mathbf{k}_1 \cdot \mathbf{k}_2 \cdots \mathbf{k}_n (\mathbb{1} + \beta^2 \mathbf{h}^2)^{ns/2} e^{-\beta |\mathbf{h}|} P$$

Since the operator $(\mathbb{1} + \beta^2 \mathbf{h}^2)^{ns/2} e^{-\beta |\mathbf{h}|}$ is bounded for any $\beta > 0, n \geq 0$, this shows that S can be written as the product of an arbitrary number of Hilbert-Schmidt operators. This implies that $S \in \mathcal{I}_p(\mathcal{K})$ for all $p > 0$ and hence $T \in \mathcal{I}_p(\mathcal{H})$ for all $p > 0$, thus proving part (ii) of the lemma. The proof of (iii) is outlined in the Appendix. Part (iv) is equivalent to the statement that the range of the map $PE_{\mathcal{C}}$ is dense in $\mathcal{H} = P\mathcal{K}$, which in turn is equivalent to the statement that the Fock state on $\mathfrak{F}$ given by P has the ‘‘Reeh-Schlieder-property’’. That this property holds in the situation under consideration has been shown in [25]. $\square$

The above lemma establishes that T has the desired property (a). It remains to be shown it also satisfies (b), i.e., that the eigenvalues t_i of T are related to the map Θ_β via eq. (45). The remainder of this proof consists in establishing this connection. For this let us first consider an arbitrary $A \in \mathfrak{F}$, and arbitrary $q_1, \dots, q_n \in \mathcal{H} = P\mathcal{K}$. Since $q_i = Pq_i$, we have that

$$\psi(q_i) = a(q_i)^* + a(PCq_i) = a(q_i)^*, \quad (49)$$

because $PCq_i = PCPq_i = P(1-P)Cq_i = 0$, using that $CP = (1-P)C$ and that $P^2 = P$. Hence, we know that

$$a(q_1)^* \dots a(q_n)^* |\Omega\rangle = \psi(q_1) \dots \psi(q_n) |\Omega\rangle, \quad (50)$$

as well as

$$\psi(q_i)^* |\Omega\rangle = a(q_i) |\Omega\rangle = 0. \quad (51)$$

Using the first relation (50), we can write

$$\langle a(q_1)^* \dots a(q_n)^* \Omega | A \Omega \rangle = \langle \Omega | \psi(q_n)^* \dots \psi(q_1)^* A | \Omega \rangle. \quad (52)$$

Applying the second relation (51), we have

$$\langle a(q_1)^* \dots a(q_n)^* \Omega | A \Omega \rangle = \langle \Omega | [\psi(q_n)^* \dots [\psi(q_2)^*, [\psi(q_1)^*, A]_\gamma]_\gamma \dots]_\gamma \Omega \rangle, \quad (53)$$

because the terms arising from the multiple (graded) commutators having one or more factor of $\psi(q_i)^*$ standing to the right of A make no contribution on account of eq. (51). We now use this relation to estimate the quantity $|\langle \Phi_{i_1 \dots i_k} | \Theta_\beta(A) \rangle|$, where $A \in \mathfrak{F}(\mathcal{O}_\mathcal{C})$, and $|\Phi_{i_1 \dots i_k}\rangle = a(p_{i_1})^* \dots a(p_{i_k})^* |\Omega\rangle$. We have

$$\begin{aligned} \langle \Phi_{i_1 \dots i_k} | \Theta_\beta(A) \rangle &= \langle a(p_{i_1})^* \dots a(p_{i_k})^* \Omega | e^{-\beta H} A \Omega \rangle = \\ &= \langle a(Pe^{-\beta \mathbf{h}} p_{i_1})^* \dots a(Pe^{-\beta \mathbf{h}} p_{i_k})^* \Omega | A \Omega \rangle, \end{aligned} \quad (54)$$

where it was used that $p_i = Pp_i$ by definition. Next, apply eq. (53) with $q_i = Pe^{-\beta \mathbf{h}} p_i$. This gives

$$\begin{aligned} \langle \Phi_{i_1 \dots i_k} | \Theta_\beta(A) \rangle &= \\ &= \langle \Omega | [\psi(Pe^{-\beta \mathbf{h}} p_{i_k})^* \dots [\psi(Pe^{-\beta \mathbf{h}} p_{i_2})^*, [\psi(Pe^{-\beta \mathbf{h}} p_{i_1})^*, A]_\gamma]_\gamma \dots]_\gamma \Omega \rangle. \end{aligned} \quad (55)$$

So far we have not used our assumption that A is localized in the region $\mathcal{O}_\mathcal{C}$, i.e., $A \in \mathfrak{F}(\mathcal{O}_\mathcal{C})$. In order to exploit this fact we write

$$Pe^{-\beta \mathbf{h}} p_i = E_\mathcal{C} Pe^{-\beta \mathbf{h}} p_i + (1 - E_\mathcal{C}) Pe^{-\beta \mathbf{h}} p_i, \quad (56)$$

where $E_{\mathcal{C}}$ denotes multiplication by the characteristic function of the subset $\mathcal{C} \subset \Sigma$. Inserting this decomposition, we have

$$\begin{aligned} [\psi(Pe^{-\beta\mathbf{h}}p_{i_1})^*, A]_{\gamma} &= [\psi((1 - E_{\mathcal{C}})Pe^{-\beta\mathbf{h}}p_{i_1})^*, A]_{\gamma} + [\psi(E_{\mathcal{C}}Pe^{-\beta\mathbf{h}}p_{i_1})^*, A]_{\gamma} \\ &= [\psi(E_{\mathcal{C}}Pe^{-\beta\mathbf{h}}p_{i_1})^*, A]_{\gamma}, \end{aligned} \quad (57)$$

where we have used in the second step that, due to graded locality (19), A has vanishing graded commutator with $\psi((1 - E_{\mathcal{C}})Pe^{-\beta\mathbf{h}}p_{i_1})^*$ because the latter is localized in the causal complement of $\mathcal{O}_{\mathcal{C}}$ by virtue of the multiplication by $1 - E_{\mathcal{C}}$, the characteristic function of $\Sigma \setminus \mathcal{C}$. Thus, on the right side of eq. (55), we are allowed to replace the expression $Pe^{-\beta\mathbf{h}}p_{i_1}$ in the innermost commutator by the localized expression $E_{\mathcal{C}}Pe^{-\beta\mathbf{h}}p_{i_1}$. We now repeat this procedure for the next-to-innermost commutator and so fourth. After k steps, we arrive at the identity

$$\begin{aligned} \langle \Phi_{i_1 \dots i_k} | \Theta_{\beta}(A) \rangle &= \\ \langle \Omega | [\psi(E_{\mathcal{C}}Pe^{-\beta\mathbf{h}}p_{i_k})^* \dots [\psi(E_{\mathcal{C}}Pe^{-\beta\mathbf{h}}p_{i_2})^*, [\psi(E_{\mathcal{C}}Pe^{-\beta\mathbf{h}}p_{i_1})^*, A]_{\gamma} \dots]_{\gamma} \Omega \rangle. \end{aligned} \quad (58)$$

We can now substitute the definition of S (see eq. (47)) into the right side of eq. (58), giving

$$\langle \Phi_{i_1 \dots i_k} | \Theta_{\beta}(A) \rangle = 2^{-k} \langle \Omega | [\psi(Sp_{i_k})^* \dots [\psi(Sp_{i_2})^*, [\psi(Sp_{i_1})^*, A]_{\gamma} \dots]_{\gamma} \Omega \rangle. \quad (59)$$

Using next that $S = VT$ and $Tp_i = t_i p_i$, we arrive at

$$\langle \Phi_{i_1 \dots i_k} | \Theta_{\beta}(A) \rangle = 2^{-k} t_{i_1} \dots t_{i_k} \langle \Omega | [\psi(Vp_{i_k})^* \dots [\psi(Vp_{i_2})^*, [\psi(Vp_{i_1})^*, A]_{\gamma} \dots]_{\gamma} \Omega \rangle. \quad (60)$$

The right side can be estimated using the Cauchy-Schwarz inequality, giving

$$|\langle \Phi_{i_1 \dots i_k} | \Theta_{\beta}(A) \rangle| \leq 2^k \cdot 2^{-k} \cdot t_{i_1} \dots t_{i_k} \|A\| \prod_{j=1}^k \|\psi(Vp_{i_j})\|, \quad (61)$$

where it was used that there are a total number of 2^k terms arising from the k repeated commutators, each of which is estimated in the same fashion by the Cauchy-Schwarz inequality. We finally use the inequalities $\|\psi(k)\| \leq \|k\|$, $k \in \mathcal{K}$, and $\|Vp_i\| \leq \|V\| \|p_i\| = 1$, since V is an isometry. This immediately gives $\|\psi(Vp_i)\| \leq 1$, and thereby the desired inequality (45). □

The proof of the theorem implies the following proposition.

Proposition IV.1. The p -norm of $\Theta_{\beta} : \mathfrak{F}(\mathcal{O}_{\mathcal{C}}) \rightarrow \mathcal{F}$ is bounded by

$$\|\Theta_{\beta}\|_p \leq [\det(\mathbf{1} + (S^*S)^{p/2})]^{1/p}, \quad \text{where } S = 2E_{\mathcal{C}}e^{-\beta\mathbf{h}}P. \quad (62)$$

Here, $E_{\mathcal{C}}$ is the projection onto the closed subspace $L^2(DM \upharpoonright \mathcal{C})$ of $\mathcal{H}$, $\mathbf{h}$ is the 1-particle Hamiltonian and P projects onto the positive spectral subspace of $\mathbf{h}$. Furthermore, we have the following estimate for $\nu_{\mathcal{C},\beta}$:

$$\nu_{\mathcal{C},\beta} = \|\Theta_{\beta}\|_1 \leq \exp \left[\left(\frac{\beta_0}{\beta} \right)^s e^{-\beta m_0/2} \right] \quad (63)$$

where $\beta_0 > 0$ is a constant depending only on the spatial geometry within $\mathcal{C}$, $m_0 = v_0 m$ and s is the number of spatial dimensions. If the spacetime metric is rescaled by $g_{ab} \rightarrow \lambda^2 g_{ab}$, then the nuclearity index for the rescaled spacetime (with Killing field $\lambda^{-1}t^a$) is estimated by the right side of (63), with β_0 replaced by $\lambda\beta_0$.

Proof. The inequality (62) is just eq. (46). The estimate for $\nu_{\mathcal{C},\beta}$ follows from this and (iii) of Lem. IV.1. In order to prove the last statement of the proposition, we note that the nuclearity index in the rescaled spacetime with Killing field $\lambda^{-1}t^a$ is equal to the nuclearity index in the unscaled spacetime at inverse temperature $\lambda^{-1}\beta$ and mass λm . Since v_0 does not change under these rescalings, this means that the nuclearity index for the rescaled spacetime is given estimated by the same expression as for the unscaled spacetime, but with β_0 replaced by $\lambda\beta_0$. $\square$

V The split-property

We are now going to show that the nuclearity property for Dirac fields in static spacetimes derived in the previous section implies the split-property.

Theorem V.1. (Split-property) Let (M, g_{ab}) be an arbitrary globally hyperbolic spacetime and ω a quasifree Hadamard state on $\mathfrak{F}$. Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be two open subsets of a Cauchy surface Σ with $\text{clo}(\mathcal{C}_1) \subset \mathcal{C}_2$ and set $\mathcal{O}_1 = \text{int}(D(\mathcal{C}_1))$ and $\mathcal{O}_2 = \text{int}(D(\mathcal{C}_2))$. Then there exists a type I factor $\mathcal{N}$ on $\mathcal{F}_{\omega}$ interpolating between the v. Neumann algebras $\pi_{\omega}(\mathfrak{F}(\mathcal{O}_1))''$ and $\pi_{\omega}(\mathfrak{F}(\mathcal{O}_2))''$ in the sense that

$$\pi_{\omega}(\mathfrak{F}(\mathcal{O}_1))'' \subset \mathcal{N} \subset \pi_{\omega}(\mathfrak{F}(\mathcal{O}_2))''. \quad (64)$$

Remark: An analogous result holds for the algebras of observables $\mathfrak{A}(\mathcal{O}) = \mathfrak{F}_+(\mathcal{O})$.

Proof. Our proof follows the chain arguments given by Verch [27, Prop. 5] in the context of a linear Klein-Gordon field, thereby using the results obtained in the previous section and some results obtained in [13].

Let us first consider the special case when (M, g_{ab}) is ultrastatic and when ω is the ground state. In that case we can define a map Θ_{β} as in eq. (44), and we know by Thm. IV.1 that this map is nuclear with a nuclearity index estimated by the bound given

in Prop. IV.1. It can be shown (see [2, Thm. 2.1]) that such a bound entails the split-property, thus proving our theorem in the case when ω is the ground state in an ultrastatic spacetime.

In order to generalize this statement from an ultrastatic spacetime to an arbitrary globally hyperbolic spacetime (with spin-structure), one now makes use of the following facts:

- (i) The ground state on an ultrastatic spacetime is of Hadamard form.
- (ii) More generally, if (M, g_{ab}) is a spacetime which is ultrastatic in an open neighborhood of some Cauchy surface Σ , then the state obtained from the “ground state construction on Σ ” is Hadamard throughout M .

Statement (i) is a corollary of the results obtained in [24]. If the Cauchy surface Σ is compact, then one can obtain the following alternative, somewhat more direct proof of (i) using a construction of Junker [15] (and using ideas of [13] to make Junker’s construction applicable to the Dirac case). We first write the projector P on the positive spectral subspace of $\mathbf{h}$ as $P = \frac{1}{2}|\mathbf{h}|^{-1}(\mathbf{h} + |\mathbf{h}|)$. The operators $\mathbf{h}$ and $|\mathbf{h}|$ can be defined for every Cauchy surface Σ perpendicular to the timelike Killing vector t^a , and can thereby act on smooth spinor fields u on M . This makes it possible rewrite the two-point function (9) of the ground state on an ultrastatic spacetime as

$$\begin{aligned}\mathcal{S}_\omega(u_1, u_2) &= \left(C \rho_\Sigma \mathcal{S} u_1 \middle| P \rho_\Sigma \mathcal{S} u_2 \right) \\ &= \frac{1}{4} \left((\mathbf{h} + |\mathbf{h}|) \rho_\Sigma \mathcal{S} u_1 \middle| |\mathbf{h}|^{-2} (\mathbf{h} + |\mathbf{h}|) \rho_\Sigma \mathcal{S} u_2 \right) \\ &= \frac{1}{4} \left(\rho_\Sigma (it^a \nabla_a + |\mathbf{h}|) \mathcal{S} u_1 \middle| |\mathbf{h}|^{-2} \rho_\Sigma (it^a \nabla_a + |\mathbf{h}|) \mathcal{S} u_2 \right),\end{aligned}\tag{65}$$

where in the third line we have used that $it^a \nabla_a \mathcal{S} u = \mathbf{h} \mathcal{S} u$ for all smooth spinor fields u on M , by the Dirac equation. The key observation is that in an ultrastatic spacetime, we have that

$$(it^a \nabla_a - |\mathbf{h}|)(it^b \nabla_b + |\mathbf{h}|)u = (\nabla^a \nabla_a - \frac{1}{4}R - m^2)u\tag{66}$$

where R is the Ricci scalar. The operator $\mathbf{h}^2$ is the self adjoint extension on $L^2(DM \upharpoonright \Sigma)$ of the elliptic differential operator $q^{ab} \nabla_a \nabla_b + m^2$, where q_{ab} is the induced (negative definite) Riemannian metric on Σ . Therefore, by standard results about powers of positive elliptic pseudo differential operators on compact manifolds (see e.g. [9]), $|\mathbf{h}|$ is a pseudo differential operator of order 1 on Σ with principal symbol $\sqrt{-q^{ab}(x)} \xi_a \xi_b I$, where I is the identity map in the fibers of DM , and $|\mathbf{h}|^{-2}$ is a pseudo differential operator of order -2 . Equation (65) therefore provides a presentation of the two-point function $\mathcal{S}_\omega$ to which Thm. 3.12 of [15] is applicable, and we conclude by that theorem that $\text{WF}(\mathcal{S}_\omega) \subset C^+$, thus showing that $\mathcal{S}_\omega$ is of Hadamard form. The above arguments do not apply as stated to

the case of non-compact Σ , since it is not guaranteed then that $|\mathbf{h}|$ is a pseudo differential operator. This can presumably be shown provided that the spatial metric q_{ab} satisfies suitable fall-off conditions at infinity, but would require a considerable effort.

Statement (ii) follows immediately from (i), together with the second remark following Def. II.1.

It is known that if the split property holds for a state ω , then it also holds for any other state that is locally quasiequivalent to ω . Therefore, since the ground state on an ultrastatic spacetime is Hadamard by item (i) of the above lemma, and since all Hadamard states are locally quasiequivalent by Thm. III.1, we conclude that the split-property holds for *any* quasifree Hadamard state on an ultrastatic spacetime. More generally, by the same argument and (ii) of the above lemma, it follows that if a spacetime has a Cauchy surface Σ with an ultrastatic neighborhood, then the split property holds for any quasifree Hadamard state and any pair of concentric double cones $\mathcal{O}_1$ and $\mathcal{O}_2$ whose bases are in Σ .

In order to further generalize this to arbitrary quasifree Hadamard states on an arbitrary globally hyperbolic spacetime, we now employ a deformation argument identical to the one given in [27, Proof of Prop. 5]. For this, one notes that one can construct, besides the original, arbitrary globally hyperbolic spacetime (M, g_{ab}) (with spin structure), a deformed spacetime $(\widehat{M}, \widehat{g}_{ab})$ with spin structure, with the following properties:

1. (M, g_{ab}) possesses a neighborhood U of some Cauchy surface Σ which is isometric to some neighborhood $\widehat{U} \subset \widehat{M}$ containing a Cauchy surface $\widehat{\Sigma}$ in the deformed spacetime $(\widehat{M}, \widehat{g}_{ab})$. We chose the spin structure of $\widehat{M}$ so that its restriction to $\widehat{U}$ coincides (via the isometry) with the restriction of spin structure of M to U ¹³.
2. The spacetime $(\widehat{M}, \widehat{g}_{ab})$ is ultrastatic in a neighborhood of a Cauchy surface $\widehat{S}$.
3. If $\mathcal{C}_1, \mathcal{C}_2 \subset \Sigma$ are sets as in the hypothesis of the theorem, then there exist open subsets $\widehat{\mathcal{V}}_1, \widehat{\mathcal{V}}_2 \subset \widehat{S}$ with compact closure such that $\text{clo}(\widehat{\mathcal{V}}_1) \subset \widehat{\mathcal{V}}_2$ and

$$\widehat{\mathcal{C}}_1 \subset \widehat{\Sigma} \cap D(\widehat{\mathcal{V}}_1), \quad \widehat{\mathcal{C}}_2 \supset \widehat{\Sigma} \cap J(\widehat{\mathcal{V}}_2),$$

where the sets $\widehat{\mathcal{C}}_i$ correspond to $\mathcal{C}_i$ and $\widehat{\Sigma}$ corresponds to Σ via the isometry postulated in 1).

Now let ω be an arbitrary quasifree Hadamard state on $\mathfrak{F}$ for the original spacetime, and let $\widehat{\omega}$ be the uniquely determined quasifree Hadamard state for the deformed spacetime, whose two-point function coincides with that of ω in $U \times U$ via the isometry identifying the

¹³It then follows that the spin structure over $\widehat{M}$ can be globally identified with the spin structure over M , since M resp. $\widehat{M}$ are topologically $\mathbb{R} \times \Sigma$ resp. $\mathbb{R} \times \widehat{\Sigma}$ and since the spin structures over Σ resp. $\widehat{\Sigma}$ have already been identified.

neighborhoods U and $\widehat{U}$ described in item 1). The trick is now to use the split-property for the deformed spacetime,

$$\pi_{\widehat{\omega}}(\widehat{\mathfrak{F}}(\text{int}(D(\widehat{\mathcal{V}}_1))))'' \subset \widehat{\mathcal{N}} \subset \pi_{\widehat{\omega}}(\widehat{\mathfrak{F}}(\text{int}(D(\widehat{\mathcal{V}}_2))))''. \quad (67)$$

(which is already known, by item 1) and 2) above and by what we have said so far) to prove the split-property in the undeformed spacetime. By item 3) above, we have that

$$\widehat{\mathcal{O}}_1 = \text{int}(D(\widehat{\mathcal{C}}_1)) \subset \text{int}(D(\widehat{\mathcal{V}}_1)), \quad \widehat{\mathcal{O}}_2 = \text{int}(D(\widehat{\mathcal{C}}_2)) \supset \text{int}(D(\widehat{\mathcal{V}}_2)),$$

therefore we have by (67) that

$$\pi_{\widehat{\omega}}(\widehat{\mathfrak{F}}(\widehat{\mathcal{O}}_1))'' \subset \widehat{\mathcal{N}} \subset \pi_{\widehat{\omega}}(\widehat{\mathfrak{F}}(\widehat{\mathcal{O}}_2))''.$$

Since $\widehat{\omega}$ agrees with ω on $U \times U$, this therefore shows the split-property on the undeformed spacetime. $\square$

Acknowledgements: This work was supported in part by MURST, INDAM-GNAMPA and NFS grant PHY00-90138 to the University of Chicago. The authors would like to thank R. Brunetti, K. Fredenhagen and E. Schroer for the invitation to the Oberwohlfach meeting “Microlocal Analysis in Quantum Field Theory”, at which part of this work was completed. S. Hollands would like to thank for the kind hospitality and financial support during his stay at the University of Rome II “Tor Vergata” from January–June 2000. He would also like to thank R. Conti and R. M. Wald for helpful discussions. C.D’Antoni thanks warmly L.Zsido and J.E.Robert for helpful discussions. We are in particular indebted to the referee, who pointed out an important shortcoming in our discussion of quasiequivalence in the original version of this paper.

A Appendix

A.1 Proof of (iii) of Lem. IV.1

We here sketch how to obtain the estimate $\|S\|_1 = \|T\|_1 \leq (\beta_0/\beta)^s e^{-\beta m_0/2}$ claimed in item (iii) of Lem. IV.1. The method of our proof also allows us to establish that the operators $\mathbf{k}_n$ introduced in the proof of Lem. IV.1 are of Hilbert-Schmidt class, as had been claimed there. However for brevity, will explicitly demonstrate this only for the case $n = 1$, the other cases can be treated in a similar way.

In order to prove the above estimate on the trace-norm of S , we write $S = \sqrt{2}E_{\mathcal{C}} \cdot \mathbf{k}_1 \cdot \mathbf{k}_2 \cdot (\mathbb{1} + \beta^2 \mathbf{h}^2)^s e^{-\beta \mathbf{h}} P$ as in the proof of Lem. IV.1, where

$$\mathbf{k}_1 = M_{\chi}(\mathbb{1} + \beta^2 \mathbf{h}^2)^{-s/2}, \quad \mathbf{k}_2 = (\mathbb{1} + \beta^2 \mathbf{h}^2)^{s/2} M_{\chi}(\mathbb{1} + \beta^2 \mathbf{h}^2)^{-s},$$

and where $M_\chi k = \chi k$ is the multiplication operator by a compactly supported smooth function χ which is identically one on $\mathcal{C}$. We get from this the estimate

$$\|S\|_1 \leq \|\mathbf{k}_1\|_2 \|\mathbf{k}_2\|_2 \|(\mathbf{1} + \beta^2 \mathbf{h}^2)^s e^{-\beta \mathbf{h}} P\| \leq c \|\mathbf{k}_1\|_2 \|\mathbf{k}_2\|_2 e^{-\beta m_0/2}.$$

Hence, in order to prove the estimate in item (iii) of Lem. IV.1, it is sufficient to demonstrate that $\|\mathbf{k}_1\|_2 \leq c\beta^{-s/2}$ and that $\|\mathbf{k}_2\|_2 \leq c\beta^{-s/2}$, where c is some constant independent of β . We will now show how to obtain the bound on the Hilbert-Schmidt norm of $\mathbf{k}_1$. The bound on the Hilbert-Schmidt norm of $\mathbf{k}_2$, and likewise on the other $\mathbf{k}_n$, can be established in a similar way, but we shall not discuss this here.

The task is thus to prove that

$$\|\mathbf{k}_1\|_2^2 = \|M_\chi(\mathbf{1} + \beta^2 \mathbf{h}^2)^{-s} M_\chi\|_1 \leq c\beta^{-s}. \quad (68)$$

Let us assume, for simplicity, that $\text{supp}(\chi) \subset \Sigma$ can be covered by a single chart $X \subset \mathbb{R}^s$, and identify $\mathcal{C}$ via this chart with a subset of X .¹⁴ The idea of the proof is to split $M_\chi(\mathbf{1} + \beta^2 \mathbf{h}^2)^{-s} M_\chi$, viewed now as a self-adjoint operator on $L^2(X; \mathbb{C}^N)$ (where $N = 2^{[(s+1)/2]}$ is the number of spinor components in $s+1$ dimensions), into two parts which are more amenable to an analysis than this operator itself. In order to achieve this, we must first recall the notion of a pseudo differential operator with a parameter (for details, see e.g. [9, p.58]). These are defined—just as ordinary pseudo differential operators—in terms of so-called symbols, the only difference being that these symbols now depend on an additional parameter, λ , which we shall take to be β^{-2} later on. Let $S_\lambda^m(X)$ be the space of all matrix valued functions $p(x, \xi, \lambda)$ which are smooth in $x \in X$ and $\xi \in \mathbb{R}^s$ and analytic in $\lambda \in \mathbb{C} \setminus \mathbb{R}^-$ and for which $|\partial_\xi^\alpha \partial_x^\gamma p| \leq c_{\alpha, \gamma} (1 + |\xi| + |\lambda|^{1/2})^{m-|\alpha|}$, where we use the usual multiindex notation, for example $\partial_x^\alpha = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \dots \frac{\partial^{\alpha_s}}{\partial x_s^{\alpha_s}}$. Elements in $S_\lambda^m(X)$ are called “symbols with a parameter”. A pseudo differential operator with parameter λ is an operator of the form

$$P_\lambda u(x) = (2\pi)^{-s/2} \int_{\mathbb{R}^s} p(x, \xi, \lambda) e^{ix\xi} \widehat{u}(\xi) d^s \xi,$$

where p is a symbol with parameter and $\widehat{u}$ is the Fourier transform of u . It is customary to write $\sigma P_\lambda = p$. The space of all pseudo differential operators P_λ with $\sigma P_\lambda \in S_\lambda^m(X)$ is denoted by $\Psi_\lambda^m(X)$.

It follows from (25) that $\mathbf{h}^2$ is a self adjoint extension of an elliptic differential operator A with symbol

$$\sigma A(x, \xi) = a_2(x, \xi) + a_1(x, \xi) + a_0(x, \xi). \quad (69)$$

The principal symbol $a_2(x, \xi)$ is given by the matrix $q^{ij}(x) \xi_i \xi_j I$ in a local coordinate chart, where I denotes the $N \times N$ identity matrix and q^{ij} are the coordinate components

¹⁴If $\mathcal{C}$ cannot be covered by single coordinate chart, one can write $\chi = \sum \chi_i$, where each χ_i is supported in a single chart. The argument then still goes through but we do not give any details here.

of the inverse spatial metric. a_1 and a_0 are symbols of order 1 and 0, respectively, whose particular form is not relevant for our purposes.

We define the symbols

$$\begin{aligned} b_0 &= (a_2 + \lambda I)^{-1} \in S_\lambda^{-2}(X), \\ b_k &= -(a_2 + \lambda I)^{-1} \sum_{2+|\alpha|+j-i=k, j < k} i^{|\alpha|} \partial_\xi^\alpha b_j \partial_x^\alpha a_i / \alpha! \in S_\lambda^{-2-k}(X). \end{aligned}$$

Let $B_\lambda \in \Psi_\lambda^{-2}(X)$ be a pseudo differential operator whose symbol has the asymptotic expansion

$$\sigma B_\lambda \sim \sum_{k \geq 0} b_k(x, \xi, \lambda) \in S_\lambda^{-2}(X). \quad (70)$$

By construction, B_λ is an inverse modulo $\Psi_\lambda^{-\infty}(X)$ of A on X (see e.g. [9]), in the sense that

$$B_\lambda(\lambda \mathbb{1} + A) = (\lambda \mathbb{1} + A)B_\lambda = \mathbb{1} \quad \text{modulo } \Psi_\lambda^{-\infty}(X). \quad (71)$$

Moreover, since A is formally self adjoint and positive, it follows from well known arguments [9] that the operator B_λ can be assumed to be a positive operator (this can be achieved, if necessary, by adding a suitable element in $\Psi_\lambda^{-\infty}(X)$ to B_λ).

We now write

$$M_\chi(\lambda \mathbb{1} + \mathbf{h}^2)^{-s} M_\chi = Q_\lambda + R_\lambda, \quad (72)$$

where $\lambda > 0$ from now on and

$$Q_\lambda = M_\chi B_\lambda^s M_\chi, \quad R_\lambda = M_\chi((\lambda \mathbb{1} + A)^{-s} - B_\lambda^s) M_\chi. \quad (73)$$

(Here and in the following we write the operator $(\lambda \mathbb{1} + \mathbf{h}^2)^{-s}$ as $(\lambda \mathbb{1} + A)^{-s}$, to simplify the notation. Note that the latter is understood as an operator on $\mathcal{K}$. This operator is not assumed to be pseudo differential, nor must it be confused with B_λ^s , which is pseudo differential, but only defined on X .)

The relevant properties of R_λ, Q_λ are summarized in the following lemma:

Lemma A.1. We have $\|Q_\lambda\|_1 \leq c\lambda^{-s/2}$ and $\|R_\lambda\|_1 \leq c\lambda^{-s/2}$ for some constant c .

The lemma allows us to finish the proof of eq. (68) because, setting $\lambda = \beta^{-2}$,

$$\|M_\chi(\mathbb{1} + \beta^2 \mathbf{h}^2)^{-s} M_\chi\|_1 \leq \beta^{-2s} (\|Q_{\beta^{-2}}\|_1 + \|R_{\beta^{-2}}\|_1) \leq \beta^{-2s} \cdot c\beta^s = c\beta^{-s}.$$

□

It remains to prove lemma (A.1).

Proof. Since Q_λ is a positive operator in $\Psi_\lambda^{-2s}(X)$, we can calculate its trace norm by (here and in the following we use the constant convention, meaning that different numerical constants, are denoted by the same symbol c):

$$\begin{aligned}
\|Q_\lambda\|_1 = \text{tr}_{L^2(X; \mathbb{C}^N)} Q_\lambda &= (2\pi)^{-s/2} \text{tr}_{\mathbb{C}^N} \int_X \int_{\mathbb{R}^s} q(x, \xi, \lambda) d^s \xi d^s x \\
&\leq c \int_{\mathbb{R}^s} (1 + |\xi| + \lambda^{1/2})^{-2s} d^s \xi \\
&\leq c \int_0^\infty (1 + r + \lambda^{1/2})^{-2s} r^{s-1} dr \\
&= c(1 + \lambda^{1/2})^{-s} \leq c\lambda^{-s/2}.
\end{aligned}$$

This proves the first inequality. For notational simplicity, we will only explicitly prove the second inequality for the case $s = 1$, but our arguments can be generalized straightforwardly to deal with general values for s . We first introduce some notation. Let $\chi', \chi'', \chi''', \dots$ be smooth functions of compact support in X such that $\chi' \equiv 1$ on the support of χ , $\chi'' \equiv 1$ on the support of χ' , $\chi''' \equiv 1$ on the support of χ'' , etc. Furthermore, let us set

$$L = [M_\chi, A], \quad L' = [M_{\chi'}, A], \quad L'' = [M_{\chi''}, A], \dots, \quad (74)$$

so that $L, L', L'', \dots$ are partial differential operators of degree 1. Let $\|\cdot\|_{H^r}$ denote the norm on the Sobolev space $H^r(X; \mathbb{C}^N)$, and let k be an arbitrary smooth spinor field. Since A is elliptic, we obtain the estimate:

$$\begin{aligned}
\|R_\lambda k\|_{H^r} &= \|M_\chi((\lambda \mathbb{1} + A)^{-1} - B_\lambda)M_\chi k\|_{H^r} \\
&\leq c\|(\lambda \mathbb{1} + A)M_\chi((\lambda \mathbb{1} + A)^{-1} - B_\lambda)M_\chi k\|_{H^{r-2}}
\end{aligned}$$

If we move the operator $(\lambda \mathbb{1} + A)$ through M_χ , then this can be further estimated by

$$\begin{aligned}
&\leq c\left(\|M_\chi(\mathbb{1} - (\lambda \mathbb{1} - A)B_\lambda)M_\chi k\|_{H^{r-2}} + \|LM_{\chi'}((\lambda \mathbb{1} - A)^{-1} - B_\lambda)M_\chi k\|_{H^{r-2}}\right) \\
&\leq c\left((1 + \lambda)^{-m/2}\|k\|_{L^2} + \|M_{\chi'}((\lambda \mathbb{1} - A)^{-1} - B_\lambda)M_\chi k\|_{H^{r-1}}\right),
\end{aligned}$$

where we have used that $\mathbb{1} - (\lambda \mathbb{1} + A)B_\lambda \in \Psi_\lambda^{-\infty}(X)$ in the last line, and where m is an arbitrary but fixed natural number. We may repeat the above chain of inequalities for the second expression in the last line, replacing χ' by χ'' and L by L' . If we iterate this

procedure, we get

$$\begin{aligned}
\|R_\lambda k\|_{H^r} &\leq c \left((1+\lambda)^{-m/2} \|k\|_{L^2} + \|M_{\chi''}((\lambda \mathbb{1} - A)^{-1} - B_\lambda) M_\chi k\|_{H^{r-2}} \right) \\
&\leq c \left((1+\lambda)^{-m/2} \|k\|_{L^2} + \|M_{\chi'''}((\lambda \mathbb{1} - A)^{-1} - B_\lambda) M_\chi k\|_{H^{r-3}} \right) \\
&\leq \dots \\
&\leq c \left((1+\lambda)^{-m/2} \|k\|_{L^2} + \|M_{\chi'''\dots'}((\lambda \mathbb{1} - A)^{-1} - B_\lambda) M_\chi k\|_{L^2} \right) \\
&\leq c(1+\lambda)^{-m/2} \|k\|_{L^2},
\end{aligned}$$

where neither of the above constants depend on λ , but only on r and m . This inequality implies that R_λ is a continuous map from $L^2(X; \mathbb{C}^N)$ to $H^r(X; \mathbb{C}^N)$ for every r . By duality, $R_\lambda^* = R_\lambda$ is a continuous map from $H^{-r}(X; \mathbb{C}^N)$ to $L^2(X; \mathbb{C}^N)$ for every r . Therefore, $K_\lambda = R_\lambda R_\lambda^*$ is a continuous map $K_\lambda : H^{-r}(X; \mathbb{C}^N) \rightarrow H^r(X; \mathbb{C}^N)$ for every r and therefore has a smooth (and by definition compactly supported) integral kernel on $X \times X$. From the dependence upon λ of the constants in the above inequality, one gets furthermore that

$$\sup |\partial_x^\alpha \partial_y^\gamma K_\lambda(x, y)| \leq c_{\alpha, \gamma} (1+\lambda)^{-m} \quad \forall \alpha, \gamma. \quad (75)$$

This estimate can now be used to bound the p -th Schatten norm $\|\cdot\|_p$ of K_λ by writing K_λ as a product of an arbitrary number of Hilbert-Schmidt operators as in eq. (42),

$$K_\lambda(x, y) = \prod_i^n (1+|x|)^{-q} \cdot (1+|x|)^{nq} K_\lambda(x, y), \quad (76)$$

where the q is chosen sufficiently large. The multiplication operator by $(1+|x|)^{-q}$ is then Hilbert-Schmidt, and $\|(1+|x|)^{nq} K_\lambda\|_2 \leq c(1+\lambda)^{-m}$ using eq. (75). Hence

$$\|K_\lambda\|_p \leq c(1+\lambda)^{-m} \quad (77)$$

for all $p > 0$ and $\lambda > 0$, where c depends only on p, m . The chain of inequalities leading to eq. (77) can be repeated for arbitrary s . It follows in particular that $\|R_\lambda\|_1^2 = \|K_\lambda\|_{1/2} \leq c(1+\lambda)^{-m} \leq c\lambda^{-s}$ (choosing $m = s$). This is the claim of the lemma for R_λ . $\square$

A.2 Nuclear maps

The notion of nuclearity considered in this paper is formulated in terms of nuclear maps. For the convenience of the reader, we now recall the definition of a nuclear map between two Banach spaces, for details we refer to [19].

Let $\mathcal{X}$ and $\mathcal{Y}$ be Banach spaces, with norms $\|\cdot\|_{\mathcal{X}}$ and $\|\cdot\|_{\mathcal{Y}}$ respectively. A bounded linear map $\Theta : \mathcal{X} \rightarrow \mathcal{Y}$ is called *nuclear*, if there exist bounded linear functionals φ_i on $\mathcal{X}$ and vectors $Y_i \in \mathcal{Y}, i \in \mathbb{I}$ such that Θ can be written as

$$\Theta(X) = \sum_i Y_i \varphi_i(X) \quad \forall X \in \mathcal{X}, \quad (78)$$

with

$$\sum_i \|\varphi_i\| \|Y_i\|_{\mathcal{Y}} < \infty, \quad \|\varphi_i\| = \sup_{\|X\|_{\mathcal{X}}=1} |\varphi_i(X)|.$$

One defines the *nuclearity index* of Θ by

$$\|\Theta\|_1 = \inf \sum_i \|\varphi_i\| \|Y_i\|_{\mathcal{Y}},$$

where the infimum is taken over all possible ways to write Θ in the form (78). The set of all nuclear maps is denoted by $\mathcal{I}_1(\mathcal{X}, \mathcal{Y})$. More generally, a bounded linear map Θ is called *p-nuclear*, if it can be written in the form (78) with

$$\|\Theta\|_p = \inf \left(\sum_i \|\varphi_i\|^p \|Y_i\|_{\mathcal{Y}}^p \right)^{1/p} < \infty, \quad p > 0.$$

The space of all *p*-nuclear maps, $p > 0$, from $\mathcal{X}$ to $\mathcal{Y}$ is denoted by $\mathcal{I}_p(\mathcal{X}, \mathcal{Y})$. It can be shown that these spaces are again Banach spaces for all $p > 0$. If $\mathcal{X} = \mathcal{Y}$, then we simply write them as $\mathcal{I}_p(\mathcal{X})$. If $\mathcal{X}$ is a Hilbert space, then $\mathcal{I}_1(\mathcal{X})$ is the space of trace-class operators, and $\mathcal{I}_2(\mathcal{X})$ is the space of Hilbert-Schmidt operators.

References

- [1] H. Araki: “On quasifree states of the CAR and Bogoliubov automorphisms,” Publ. RIMS Kyoto Univ. **6**, 385-442 (1970)
- [2] D. Buchholz, C. D’Antoni and K. Fredenhagen: “The universal structure of local algebras,” Commun. Math. Phys. **111** 123 (1987)
- [3] D. Buchholz, C. D’Antoni and R. Longo: “Nuclear Maps and Modular Structures II: Applications to Quantum Field Theory,” Commun. Math. Phys. **129**, 115-138 (1990)
- [4] D. Buchholz and E. H. Wichmann: “Causal independence and the energy-level density of states in quantum field theory,” Commun. Math. Phys. **106**, 321-344 (1986)
- [5] D. Buchholz and M. Poppmann: “How small is the phase space in quantum field theory?” Ann. Inst. H. Poincaré **52**, 237 (1991)
- [6] J. Dimock: “Dirac quantum fields on a manifold,” Trans. Am. Math. Soc. **269**, 133-147 (1982)
- [7] J. J. Foit: “Abstract Twisted Duality for Quantum Free Fermi Field,” Publ. R.I.M.S. Kyoto Univ. **19** (1983) 729-741

- [8] S. A. Fulling, F. J. Narcowich and R. M. Wald: “Singularity Structure Of The Two Point Function In Quantum Field Theory In Curved Space-Time. II,” *Annals Phys.* **136** (1981) 243.
- [9] P. B. Gilkey: “The Index Theorem and the Heat Equation,” Publish or Perish, Boston (1974)
- [10] R. Haag: “Local Quantum Physics,” Springer Verlag, Berlin (1990)
- [11] R. Haag and J. A. Swieca: “When does a quantum field theory describe particles?” *Commun. Math. Phys.* **5**, 215 (1967)
- [12] L. Hörmander: “The analysis of linear partial differential operators, III,” Springer-Verlag, Berlin (1985)
- [13] S. Hollands: “The Hadamard Condition for Dirac fields and Adiabatic States on Robertson-Walker Spacetimes,” *Commun. Math. Phys.* **216**, 635 (2001)
- [14] S. Hollands and W. Ruan: “The state space of perturbative quantum field theory in curved space-times,” *Annales Henri Poincare* **3**, 635 (2002) [arXiv:gr-qc/0108032].
- [15] W. Junker: “Hadamard States, Adiabatic Vacua and the Construction of Physical States for Scalar Quantum Fields on Curved Spacetime,” *Rev. Math. Phys.* **8** 1091-1159 (1996)
- [16] W. Junker and E. Schroer: “Adiabatic vacuum states on general spacetime manifolds: Definition, construction and physical properties,” *Annales Poincare Phys. Theor.* **3**, 1113 (2002) [arXiv:math-ph/0109010].
- [17] B. S. Kay and R. M. Wald: “Theorems On The Uniqueness And Thermal Properties Of Stationary, Nonsingular, Quasifree States On Space-Times With A Bifurcate Killing Horizon,” *Phys. Rept.* **207** (1991) 49.
- [18] K. Kratzert: “Singularity structure of the two point function of the free Dirac field on a globally hyperbolic spacetime,” *Annalen Phys.* **9** (2000) 475 [arXiv:math-ph/0003015].
- [19] A. Pietsch: “Nuclear locally convex spaces,” Springer-Verlag, Berlin (1972)
- [20] R. T. Powers and E. Størmer: “Free states of the canonical anticommutation relations,” *Commun. Math. Phys.* **16**, 1 (1970)
- [21] M. J. Radzikowski: “Micro-Local Approach To The Hadamard Condition In Quantum Field Theory On Curved Space-Time,” *Commun. Math. Phys.* **179** (1996) 529.

- [22] M. Reed and B. Simon: “Methods of modern mathematical physics, IV,” Academic Press, London (1978)
- [23] H. Sahlmann and R. Verch: “Microlocal spectrum condition and Hadamard form for vector-valued quantum fields in curved spacetime,” *Rev. Math. Phys.* **13** (2001) 1203 [arXiv:math-ph/0008029]
- [24] H. Sahlmann and R. Verch, “Passivity and microlocal spectrum condition,” *Commun. Math. Phys.* **214** (2000) 705 [arXiv:math-ph/0002021]
- [25] A. Strohmaier: “The Reeh-Schlieder property for quantum fields on stationary spacetimes,” [math-ph/0002054]
- [26] S. Summers: “Normal product states for fermions and twisted duality for CCR and CAR-type algebras with application to the Yukawa₂ quantum field model,” *Commun. Math. Phys.* **86**, 111-141 (1982)
- [27] R. Verch: “Nuclearity, split-property and duality for the Klein-Gordon field in curved space-time,” *Lett. Math. Phys.* **29**, 297 (1993)
- [28] R. Verch: “Local definiteness, primarity and quasiequivalence of quasifree Hadamard quantum states in curved space-time,” *Commun. Math. Phys.* **160**, 507 (1994)
- [29] R. Verch: “A Spin-Statistics Theorem for Quantum Fields on Curved Spacetime Manifolds in a Generally Covariant Framework,” *Commun. Math. Phys.* **223**, 261 (2001) [arXiv:math-ph/0102035].
- [30] R. Verch: “Scaling Analysis and Ultraviolet Behavior of Quantum Field Theories in Curved Spacetime,” PhD thesis, Hamburg University (1994)
- [31] S. Hollands and R. M. Wald: “Local Wick Polynomials and Local Time Ordered Products of Quantum Fields in Curved Spacetime,” *Commun. Math. Phys.* **223**, 289 (2001) [arXiv:gr-qc/0103074].
- [32] R. M. Wald: “General Relativity,” The University of Chicago Press, Chicago (1984)